

Recombinant Mouse Pdgfa Protein

Cat. No. Pdgfa-048M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant mouse platelet-derived growth factor AA (rmPDGF-AA) Ser87-Arg196(Accession # NP_032834.1) was expressed in human 293 cells (HEK293).
Species	Mouse
Source	HEK293
ProteinLength	87-196 a.a.
Description	<p>PDGFs are mitogenic during early developmental stages, driving the proliferation of undifferentiated mesenchyme and some progenitor populations. During later maturation stages, PDGF signalling has been implicated in tissue remodelling and cellular differentiation, and in inductive events involved in patterning and morphogenesis. In addition to driving mesenchymal proliferation, PDGFs have been shown to direct the migration, differentiation and function of a variety of specialised mesenchymal and migratory cell types, both during development and in the adult animal. Other growth factors in this family include vascular endothelial growth factors B and C (VEGF-B, VEGF-C) which are active in angiogenesis and endothelial cell growth, and placenta growth factor (PlGF) which is also active in angiogenesis. PDGF plays a role in embryonic development, cell proliferation, cell migration, and angiogenesis. PDGF is a required element in cellular division for fibroblast, a type of connective tissue cell. PDGF is also known to maintain proliferation of oligodendrocyte progenitor cells. Platelet-derived growth factor subunit A is also known as PDGFA, PDGF-A, PDGF1, is a member of the platelet-derived growth</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

factor family. PDGFA can exist either as a homodimer (PDGF-AA) or as a heterodimer with the platelet-derived growth factor Bata polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds.

Molecular Mass

rmPDGF-AA, with two amino acids (Gly-Pro) at the N- terminus is a homodimeric, glycosilated polypeptide chain containing 2x110 amino acids, and has a calculated MW of 2x12.7 kDa. DTT-reduced protein migrates as a 16.5 kDa polypeptide and the non-reduced cystine-linked homodimer migrates as a 28 kDa protein.

Endotoxin

Less than 1.0 EU per 1 g of the rmPDGF-AA by the LAL method.

Purity

>95% as determined by SDS-PAGE of 6 g reduced (+) and non-reduced (-) rmPDGF-AA. All lots are greater than 95% pure.

Notes

Research Use Only. Not for use in diagnostic or therapeutic procedures.

Storage

Store at -20 centigrade in lyophilized state after receipt. For long term storage, upon reconstitution rmPDGF-AA should be aliquot and store at -20 centigrade or -8 centigrade centigrade. Avoid repeated freeze-thaw cycles.

Storage Buffer

Bulk protein in a 0.22 m filtered solution of 50mM Tris 150mM NaCl pH 7.0 and delivered as liquid formulation or lyophilized powder. Normally 5-8% trehalose and mannitol are added as protectants before lyophilization.

GENE INFORMATION

Gene Name

[Pdgfa platelet derived growth factor, alpha \[Mus musculus \(house mouse\) \]](#)

Official Symbol

[Pdgfa](#)

Synonyms

Pdgfa platelet derived growth factor, alpha; PDGFA; PDGF-A; PDGF1; PDGF-1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



platelet-derived growth factor subunit A; PDGF A-chain
PDGF subunit A; platelet-derived growth factor A-chain; platelet-derived growth factor
alpha polypeptide

Gene ID 18590

mRNA Refseq NM_008808

Protein Refseq NP_032834

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA