

Active Native Bovine Prothrombin Frag 1

Cat. No. Prothrombin-271B Lot. No. (See product label)

SPECIFICATION

Species Bovine

Source Plasma

Description

Prothrombin is a vitamin K-dependent plasma protein which is synthesized in the liver. Prior to secretion into plasma, prothrombin undergoes post-translational modification by a vitamin K-dependent carboxylase which converts ten specific glutamic acid residues to γ -carboxyglutamic acid (gla). The ten gla residues are located within the first 40 amino acids of the mature protein and contribute to the ability of prothrombin to bind to negatively charged phospholipid membranes. Prothrombin contains two regions of internal homology which are referred to as "kringle" structures. These regions of conspicuous secondary structure are located between residues 40 and 270 of the mature plasma protein and replace the growth factor domains found in several other plasma serine proteases. Thus far, no function has been ascribed to these regions, but there is suspicion that they may play a role in one of several binary protein interactions involving prothrombin. The mature single chain protein circulates in plasma as a zymogen and, during coagulation, is proteolytically activated to the potent serine protease α -thrombin. This proteolysis is catalyzed by the prothrombinase enzyme complex. During activation, prothrombin is cleaved at Arg271-Thr272 (human) / Arg273-Thr274 (bovine) and at Arg320-Ser321 (human) / Arg323-Ser324 (bovine) to a "pro" fragment (fragment 1.2) and thrombin, the latter of which is composed of two chains covalently linked by a disulfide bond. In the case of human prothrombin/thrombin, there is an additional thrombin feed-back cleavage at Arg284-Thr285 resulting in an additional 13 amino acids being removed

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	from the mature thrombin "A" chain.
Form	50% (vol/vol) glycerol/H ₂ O
Bio-activity	Clotting assay
Molecular Mass	21700
Purity	>95% by SDS-PAGE. NOT tissue/cell culture grade. Not tested for endotoxin.
Characteristic	Extinction coefficient: 14.4, Plasma concentration:100 µg/ml, Isoelectric point:4.4-4.9, Percent carbohydrate:0.1, Structure: single chain, NH ₂ -terminal gla domain, two kringle regions
Storage	-20°C

GENE INFORMATION

Gene Name	F2 coagulation factor II (thrombin) [Bos taurus]
Official Symbol	F2
Synonyms	prothrombin; thrombin
Gene ID	280685
mRNA Refseq	NM_173877
Protein Refseq	NP_776302
UniProt ID	P00735

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	chromosome: 15
Pathway	Cell surface interactions at the vascular wall, organism-specific biosystem; Complement and Coagulation Cascades, organism-specific biosystem; G alpha (q) signalling events, organism-specific biosystem
Function	calcium ion binding; fibrinogen binding; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA