

Recombinant Human RBBP4 cell lysate

Cat. No. RBBP4-650HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human RBBP4 / RBAP48 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[RBBP4 retinoblastoma binding protein 4 \[Homo sapiens \]](#)

Official Symbol

RBBP4

Synonyms

RBBP4; retinoblastoma binding protein 4; histone-binding protein RBBP4; NURF55; RbAp48; RBBP-4; CAF-I p48; CAF-1 subunit C; CAF-I 48 kDa subunit; MSI1 protein homolog; retinoblastoma-binding protein 4; retinoblastoma-binding protein p48; chromatin assembly factor 1 subunit C; chromatin assembly factor I p48 subunit; chromatin assembly factor/CAF-1 p48 subunit; nucleosome-remodeling factor subunit RBAP48; RBAP48;

Gene ID

[5928](#)

mRNA Refseq

[NM_001135255](#)

Protein Refseq

[NP_001128727](#)

MIM

[602923](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q09028
Chromosome Location	1p35.1
Pathway	Cell Cycle, organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem; Chromosome Maintenance, organism-specific biosystem; Deposition of New CENPA-containing Nucleosomes at the Centromere, organism-specific biosystem; E2F transcription factor network, organism-specific biosystem; G0 and Early G1, organism-specific biosystem; Hedgehog signaling events mediated by Gli proteins, organism-specific biosystem;
Function	contributes_to DNA-dependent ATPase activity; histone binding; histone deacetylase binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA