

## Recombinant Human RELA 293 Cell Lysate

**Cat. No.** RELA-2423HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for v-rel reticuloendotheliosis viral oncogene homolog A (avian) (RELA), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	RELA v-rel reticuloendotheliosis viral oncogene homolog A (avian) [ Homo sapiens ]
<b>Official Symbol</b>	RELA
<b>Synonyms</b>	RELA; v-rel reticuloendotheliosis viral oncogene homolog A (avian); NFKB3, nuclear factor of kappa light polypeptide gene enhancer in B cells 3; transcription factor p65; p65; NF-kappa-B p65delta3; nuclear factor NF-kappa-B p65 subunit; nuclear factor of kappa light polypeptide gene enhancer in B-cells 3; NFKB3; MGC131774;
<b>Gene ID</b>	5970
<b>mRNA Refseq</b>	NM_001145138
<b>Protein Refseq</b>	NP_001138610
<b>MIM</b>	164014
<b>UniProt ID</b>	Q04206
<b>Chromosome Location</b>	11q13
<b>Pathway</b>	Activated TLR4 signalling, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Acute myeloid leukemia, organism-specific biosystem; Acute myeloid leukemia, conserved biosystem; Adaptive Immune System, organism-specific biosystem; Adipocytokine signaling pathway, organism-specific biosystem; Adipocytokine signaling pathway, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Function**

DNA binding; NF-kappaB binding; activating transcription factor binding; ankyrin repeat binding; chromatin binding; identical protein binding; identical protein binding; phosphate ion binding; protein N-terminus binding; protein binding; protein kinase binding; repressing transcription factor binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; sequence-specific DNA binding transcription factor activity; sequence-specific distal enhancer binding RNA polymerase II transcription factor activity; transcription factor binding; transcription regulatory region DNA binding; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA