

Recombinant Full Length Escherichia Coli Potassium-Transporting Atpase C Chain(Kdpc) Protein, His-Tagged

Cat. No. RFL33641EF Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Escherichia coli Potassium-transporting ATPase C chain(kdpC) Protein (B6HYQ4) (1-190aa), fused to N-terminal His tag, was expressed in E. coli.
Species	E.coli
Source	E.coli
ProteinLength	Full Length (1-190)
Form	Lyophilized powder
AA Sequence	MRGLRPALSTFIFLLLITGGVYPLLTTLVGLQWWFPWQANGSLIREGDTVIRGSALIGQ NFT GNGYFQGRPSATAEMPYPNPQASGGSNLAVSNPELDKQIAARVAALRAANPDA STNVPVEL VTASASGLDNNITPQAAVWQIPRVAKARNLSVEQLTQLIAKYSQQPLVK YIGQPVVNIVE LNLALDKLDE
Purity	Greater than 90% as determined by SDS-PAGE.
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Buffer Tris/PBS-based buffer, 6% Trehalose, pH 8.0

Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

GENE INFORMATION

Gene Name kdpC

Synonyms kdpC; ECSE_0756; Potassium-transporting ATPase KdpC subunit; ATP phosphohydrolase [potassium-transporting] C chain; Potassium-binding and translocating subunit C; Potassium-translocating ATPase C chain

UniProt ID [B6HYQ4](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA