

Recombinant Human RIPK3 293 Cell Lysate

Cat. No. RIPK3-2332HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for receptor-interacting serine-threonine kinase 3 (RIPK3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	RIPK3 receptor-interacting serine-threonine kinase 3 [Homo sapiens]
Official Symbol	RIPK3
Synonyms	RIPK3; receptor-interacting serine-threonine kinase 3; receptor-interacting serine/threonine-protein kinase 3; RIP3; RIP-3; RIP-like protein kinase 3; receptor interacting protein 3; receptor-interacting protein 3;
Gene ID	11035
mRNA Refseq	NM_006871
Protein Refseq	NP_006862
MIM	605817
UniProt ID	Q9Y572
Chromosome Location	14q12
Pathway	Cytosolic DNA-sensing pathway, organism-specific biosystem; Cytosolic DNA-sensing pathway, conserved biosystem; Cytosolic sensors of pathogen-associated DNA, organism-specific biosystem; DAI mediated induction of type I IFNs, organism-specific biosystem; Immune System, organism-specific biosystem; Innate Immune System, organism-specific biosystem; RIP-mediated NFkB activation via DAI, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

ATP binding; NF-kappaB-inducing kinase activity; nucleotide binding; protein binding; protein kinase activity; protein serine/threonine kinase activity; transcription coactivator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA