

Recombinant Human SCARB1 cell lysate

Cat. No. SCARB1-2699HCL Lot. No. (See product label)

SPECIFICATION

| | |
|--------------------------------|---|
| Product Overview | Human SCARB1 / CD36L1 / CLA-1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls |
| Species | Human |
| Source | Human Cells |
| Preparation method | Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| Lysis buffer | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF |
| Quality control Testing | 12.5% SDS-PAGE Stained with Coomassie Blue |
| Recommended Usage | 1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[SCARB1 scavenger receptor class B, member 1 \[Homo sapiens \]](#)

Official Symbol

SCARB1

Synonyms

SCARB1; scavenger receptor class B, member 1; CD36 antigen (collagen type I receptor, thrombospondin receptor) like 1 , CD36L1; scavenger receptor class B member 1; CLA 1; CLA1; SR BI; SRB1; CD36 and LIMPII analogous 1; scavenger receptor class B type III; CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 1; CLA-1; SR-BI; CD36L1; HDLQTL6; MGC138242;

Gene ID

[949](#)

mRNA Refseq

[NM_001082959](#)

Protein Refseq

[NP_001076428](#)

MIM

[601040](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|----------------------------|---|
| UniProt ID | Q8WTV0 |
| Chromosome Location | 12q24.32 |
| Pathway | Bile secretion, organism-specific biosystem; Bile secretion, conserved biosystem; Fat digestion and absorption, organism-specific biosystem; Fat digestion and absorption, conserved biosystem; Folate Metabolism, organism-specific biosystem; HDL-mediated lipid transport, organism-specific biosystem; Hepatitis C, organism-specific biosystem; |
| Function | 1-phosphatidylinositol binding; apolipoprotein A-I binding; apolipoprotein binding; high-density lipoprotein particle binding; high-density lipoprotein particle receptor activity; lipopolysaccharide binding; lipopolysaccharide receptor activity; low-density lipoprotein particle binding; phosphatidylserine binding; protein homodimerization activity; receptor activity; transporter activity; |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA