

## Active Recombinant Mouse SDC2 Protein

Cat. No. SDC2-14790M Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Recombinant mouse Syndecan-2 (Glu19-Phe141) protein with C-terminal TGHHH HHHHHGGQ was expressed in HEK293 cell.

**Species** Mouse

**Source** HEK293

**Description** Syndecan-2, previously known as fibroglycan, is a member of the syndecan family of Type 1 transmembrane proteins carrying heparan sulfate and chondroitin sulfate glycosaminoglycans. The syndecan family consists of four transmembrane proteoglycans divided into two subfamilies (Syndecan -1 , -3 and Syndecan-2 , - 4) based on their sequence homology. The four syndecans have conserved cytoplasmic domains which enables them to associate with cytoskeletal proteins and signaling molecules. The cytoplasmic tail of Syndecan-2 has both serine and tyrosine phosphorylation sites. Non-covalent homodimerization of Syndecan-2 is dependent on the transmembrane domain. Syndecan-2 is expressed in endothelial, mesenchymal, carcinoma cells, and is the predominant syndecan expressed during embryonic development. It functions as a co-receptor for enzymes, growth factors, and cytokines, and contributes to the formation of cytokine gradients through extracellular shedding. Syndecan-2 interacts with high affinity receptors and integrins to modulate intracellular signaling. Syndecan is also involved in matrix assembly and remodeling. It controls laminin and fibronectin matrix assembly into a fibrillar matrix.

**Bio-activity** When mouse Syndecan-2 is immobilized at 5 µg/mL, human FGF-basic binds with

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	EC50 of 2 - 8 ng/mL.
<b>Molecular Mass</b>	14.9 kDa
<b>N-terminal Sequence Analysis</b>	Glu
<b>Endotoxin</b>	Less than 0.1 EU per µg of protein as determined by the LAL method.
<b>Purity</b>	>95% as determined by Coomassie stained SDS-PAGE.
<b>Applications</b>	Bioassay
<b>Storage</b>	Unopened vial can be stored between 2 centigrade and 8 centigrade for one month, at -20 centigrade for six months, or at -70 centigrade for one year. For maximum results, quick spin vial prior to opening. The protein can be aliquoted and stored at -20 centigrade to -70 centigrade. Stock solutions can also be prepared at 50 - 100 µg/mL in sterile buffer (PBS, HPBS, DPBS, or EBSS) containing carrier protein such as 0.2-1% BSA or HSA and stored in working aliquots at -20 centigrade to -70 centigrade. Avoid repeated freeze/thaw cycles.
<b>Concentration</b>	200 µg/mL
<b>Storage Buffer</b>	0.22 µm filtered protein solution is in PBS, pH 7.2.
<b>GENE INFORMATION</b>	
<b>Gene Name</b>	<a href="#">Sdc2 syndecan 2 [ Mus musculus (house mouse) ]</a>
<b>Official Symbol</b>	<a href="#">Sdc2</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>Synonyms</b>	SDC2; Synd2; Hspg1; Fibroglycan; Hspg1; Synd2; AA960457; syndecan-2; 4833414L08Rik; syndecan-2; HSPG; fibroglycan; heparan sulfate proteoglycan 1, cell surface-associated (fibroglycan); heparan sulfate proteoglycan core protein
<b>Gene ID</b>	<a href="#">15529</a>
<b>mRNA Refseq</b>	<a href="#">NM_008304</a>
<b>Protein Refseq</b>	<a href="#">NP_032330</a>
<b>UniProt ID</b>	<a href="#">P43407</a>

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA