

## Recombinant Human SERPINB2

Cat. No. SERPINB2-1554H Lot. No. (See product label)

### SPECIFICATION

#### Product Overview

Recombinant Human serpin peptidase inhibitor, clade B (ovalbumin), member 2 was expressed in *E. coli* Cells. MW = 47kDa.

#### Species

Human

#### Source

E.coli

#### Description

PAI-2 is an inhibitory serpin expressed mainly in keratinocytes, activated monocytes, and placental trophoblasts. It exists predominantly as a 47 kDa nonglycosylated intracellular protein which can be induced to be secreted as 60 kDa glycoprotein. The physiological relevance of this property, which is neither a consequence of any mutation in the PAI-2 gene nor associated with any known disorder, is still unclear. However, it appears that the formation of intracellular dormant polymers may be important for the controlled release of the inhibitor from PAI-2 producing cells. Plasma levels of PAI-2 are usually low or undetectable, except during pregnancy and in some forms of monocytic leukemia. Secretion of PAI-2 from the placenta normally occurs during the third trimester of pregnancy and accounts for the dramatic increase in PAI-2 levels (up to 250 ng/ml), which are maintained at these levels until postpartum, and then rapidly decline. In addition to its vital role in protecting the placenta from degradation by uPA and/or uPA-activated proteases, PAI-2 has been shown to be essential for the prevention of metastatic spread of neck, lung and breast cancers. The beneficial effect of PAI-2 seen in these studies is presumed to stem from its ability to inhibit uPA-dependent cell dissemination. PAI-2 has also been reported to inhibit keratinocyte proliferation, and to participate in the innate immune response

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

during viral infection. Recombinant human PAI-2 is a 415-residue nonglycosylated protein.

**Sequence**

MEDLCVANTL FALNLFKHLA KASPTQNLFL SPWSISSTMA MVYMGSRGST  
 EDQMAKVLQF NEVGANAVT PMTPENFTSC GFMQQIQKGS YPDAILQAQA  
 ADKIHSSFRS LSSAINASTGN YLLESVNK LFGEKSASFR EEYIRLCQKY  
 YSSEPQAVDF LECAEEARKK INSWVKTQTK GK IPNLLPE GSVGDTRMV  
 LVNAVYFKGK WKTPFEKKN GLYPFRVNSA QRT.

**Purity**

Greater than 95% by SDS-PAGE gel and HPLC analyses. Endotoxin level is less than 0.1 ng per µg (1EU/µg).

**Storage**

The lyophilized PAI-2 recombinant protein is stable for at least 2 years from date of receipt at -20°C. Reconstituted PAI-2 is stable for at least 3 months when stored in working aliquots with a carrier protein at -20°C. As with any protein, exposing PAI-2 recombinant protein to repeated freeze / thaw cycles is not recommended. When working with proteins care should be taken to keep recombinant protein at a cool and stable temperature.

## GENE INFORMATION

**Gene Name** [SERPINB2 serpin peptidase inhibitor, clade B \(ovalbumin\), member 2 \[ Homo sapiens \]](#)

**Synonyms** SERPINB2; serpin peptidase inhibitor, clade B (ovalbumin), member 2; PAI; PAI2; PAI-2; PLANH2; HsT1201; serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2; Plasminogen activator inhibitor 2; Plasminogen activator inhibitor 2; Monocyte Arg-serpin; Monocyte Arg-serpin

**Gene ID** [5055](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>mRNA Refseq</b>	NM_001143818
<b>Protein Refseq</b>	NP_001137290
<b>MIM</b>	173390
<b>UniProt ID</b>	P05120
<b>Chromosome Location</b>	18q21.3
<b>Pathway</b>	Hemostasis
<b>Function</b>	peptidase inhibitor activity; serine-type endopeptidase activity; serine-type endopeptidase inhibitor activity

PDB rendering based on 1by7.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA