

Active Recombinant Human SERPINB2 protein, His-tagged

Cat. No. SERPINB2-5060H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human SERPINB2 protein(P05120)(1~415 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	1-415 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Serpin B2, also known as PAI-2, is an approximately 60 kDa serine protease inhibitor. It is primarily secreted by macrophages and monocytes and can form disulfide-linked multimers. Serpin B2 inhibits both the urokinase-type and tissue-type plasminogen activators (uPA and tPA). Serpin B2 also promotes the clearance of uPA by enhancing its binding and uptake by LRP. It limits fibril formation by Huntington protein (HTT) and beta-Amyloid peptides. It promotes Th2 biased immune responses and is important for intestinal CCL2 production, monocyte recruitment, and nematode clearance. A non-glycosylated form of Serpin B2 is retained intracellularly where it interferes with TNF- α induced apoptosis by protecting the Retinoblastoma protein (RB1) from calpain digestion. It also inhibits proteasome activity in activated endothelial cells. The activity of recombinant human PAI-2 was measured by its ability to inhibit uPA cleavage of a peptide substrate, N-carbobenzyloxy-Gly-Gly-Arg-7-amido-4-methylcoumarin (Z-GGR-AMC)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

in the assay buffer 50 mM Tris, 0.01% Tween 20, pH 8.5. The 50 ul different concentrations of rhPAI-2 (MW: 48.2 KD) was incubated with 50ul 2ug/ml rhuPA (EPA140Mu6 1) at room temperature for 15 minutes. Loading 50 μ of the incubated mixtures into empty wells of a plate, and start the reaction by adding 50 μ of 200 μM substrate (Z-GG R-AMC). Include a substrate blank containing 50 μ of assay buffer and 50 μ of 200 μM substrate. Then read at excitation and emission wavelengths of 380 nm and 460 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that recombinant human PAI2 significantly decreased uPA activity. The inhibition IC50 was <28 nM.

Molecular Mass 48.2kDa

Endotoxin <1.0EU per 1μg (determined by the LAL method).

Purity > 80% as determined by SDS-PAGE.

Storage Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name SERPINB2 serpin peptidase inhibitor, clade B (ovalbumin), member 2 [Homo sapiens]

Official Symbol SERPINB2

Synonyms SERPINB2; serpin peptidase inhibitor, clade B (ovalbumin), member 2; PAI2, PLANH2, serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

plasminogen activator inhibitor 2; HsT1201; serpin B2; monocyte Arg-serpin; urokinase inhibitor; placental plasminogen activator inhibitor; plasminogen activator inhibitor, type II (arginine-serpin); serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2; PAI; PAI2; PAI-2; PLANH2;

Gene ID [5055](#)


mRNA Refseq [NM_001143818](#)

Protein Refseq [NP_001137290](#)

MIM [173390](#)

UniProt ID [P05120](#)

SDS-PAGE




 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA