

Recombinant Human SETD8 293 Cell Lysate

Cat. No. SETD8-1924HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for SET domain containing (lysine methyltransferase) 8 (SETD8) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	SETD8 SET domain containing (lysine methyltransferase) 8 [Homo sapiens]
Official Symbol	SETD8
Synonyms	SETD8; SET domain containing (lysine methyltransferase) 8; N-lysine methyltransferase SETD8; KMT5A; PR Set7; SET07; SET8; PR/SET07; H4-K20-HMTase SETD8; lysine N-methyltransferase 5A; SET domain-containing protein 8; PR/SET domain containing protein 8; PR/SET domain-containing protein 07; histone-lysine N-methyltransferase SETD8; H4-K20-specific histone methyltransferase; H4K20-specific histone methyltransferase splice variant Set8b; PR-Set7;
Gene ID	387893
mRNA Refseq	NM_020382
Protein Refseq	NP_065115
MIM	607240
UniProt ID	Q9NQR1
Chromosome Location	12q24.31
Pathway	Lysine degradation, organism-specific biosystem; Lysine degradation, conserved biosystem; p53 pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

histone-lysine N-methyltransferase activity; methyltransferase activity; p53 binding; protein binding; protein-lysine N-methyltransferase activity; transcription corepressor activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA