

Recombinant Human SFPQ 293 Cell Lysate

Cat. No. SFPQ-1911HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for splicing factor proline/glutamine-rich (polypyrimidine tract binding protein associated) (SFPQ) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name SFPQ splicing factor proline/glutamine-rich [Homo sapiens]

Official Symbol SFPQ

Synonyms SFPQ; splicing factor proline/glutamine-rich; splicing factor proline/glutamine rich (polypyrimidine tract binding protein associated); splicing factor, proline- and glutamine-rich; polypyrimidine tract binding protein associated; PSF; hPOMp100; 100 kDa DNA-pairing protein; PTB-associated splicing factor; PTB-associated-splicing factor; DNA-binding p52/p100 complex, 100 kDa subunit; polypyrimidine tract-binding protein-associated splicing factor; polypyrimidine tract-binding protein-associated-splicing factor; splicing factor proline/glutamine rich (polypyrimidine tract-binding protein-associated); POMP100; DKFZp547C228; DKFZp667K0521; DKFZp686K1282;

Gene ID 6421

mRNA Refseq NM_005066

Protein Refseq NP_005057

MIM 605199

UniProt ID P23246

Chromosome Location 1p34.3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Pathway	mRNA processing, organism-specific biosystem;
Function	DNA binding; RNA binding; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA