

## Active Recombinant Human SIRPA protein, Fc-tagged

**Cat. No.** SIRPA-202H    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human SIRPA(Glu31-Arg370) fused with Fc region of Human IgG1 at C-terminal was expressed in CHO.
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	31-370 a.a.
<b>Description</b>	<p>Signal regulatory protein alpha (SIRP alpha, designated CD172a), also called SHPS-1 (SHP substrate 1) and previously, MyD-1 (Myeloid/Dendritic-1), is a monomeric ~90 kDa type I transmembrane glycoprotein that belongs to the SIRP/SHPS (CD172) family of the immunoglobulin superfamily. SIRPs are paired receptors, with similar extracellular domains but differing C-termini and functions. The 503 amino acid (aa) human SIRP alpha contains a 342 aa extracellular domain (ECD), with one V-type, and two C1 type Ig domains, and three potential N glycosylation sites. It has a 110 aa cytoplasmic sequence with ITIM motifs that recruit tyrosine phosphatases SHP-1 and SHP-2 when phosphorylated. Human SIRP alpha has more than 40 described polymorphisms, including the prominent BIT (Brain Ig like molecule with Tyrosine-based activation motifs, also called SIRP alpha 2 or PTPNS. One reported isoform lacks aa 1-101, which eliminates most of the V type Ig domain. Human SIRP alpha ECD shares 61%, 60%, 71%, 72% and 73% aa identity with mouse, rat, porcine, bovine and equine SIRP alpha, respectively; it shares 84% and 76% aa identity with human SIRP beta 1 and SIRP gamma, respectively. SIRP alpha is expressed mainly</p>

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

on myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It is also found on neurons, smooth muscle and endothelial cells. SIRP alpha shows adhesion to the ubiquitous CD47/IAP (integrin associated protein), while SIRP gamma binds more weakly and SIRP alpha 1 does not bind at all. Mouse and human SIRP alpha -CD47 binding only cross-reacts for specific polymorphisms and influences engraftment of xenotransplanted stem cells. SIRP alpha engagement generally produces a negative regulatory signal. Low SIRP alpha recognition of CD47, which occurs on aged erythrocytes or platelets or xenogenic cells, promotes clearance of CD47<sup>low</sup> cells from circulation. SIRP alpha recognition of surfactants SP-A and SP-D in the lung can inhibit alveolar macrophage cytokine production. The CD47 integrin-SIRP alpha interaction is reported to promote macrophage fusion during osteoclastogenesis.

**Predicted N Terminal** Gly27 & Glu31

**Form** Lyophilized from a 0.2 µm filtered solution in PBS.

**Bio-activity** Measured by the ability of the immobilized protein to support the adhesion of human red blood cells. Subramanian, S. et al. (2006) Blood 107:2548. Red blood cells, added at  $10 \times 10^6$  cells/mL (100 µ/well) to SIRP alpha coated plates, will adhere after 1 hour at 37 centigrade. The ED50 for this effect is 0.6-2.4 g/mL.

**Molecular Mass** Predicted Molecular Mass: 64 (unlabeled) kDa

**Endotoxin** <0.1 EU per 1 µg of the protein by the LAL method.

**Purity** >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

**Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

12 months from date of receipt, -20 to -70 centigrade as supplied.  
 1 month, 2 to 8 centigrade under sterile conditions after reconstitution.  
 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

**Reconstitution** Reconstitute at 500 µg/mL in PBS.

## GENE INFORMATION

**Gene Name** [SIRPA signal-regulatory protein alpha \[ Homo sapiens \]](#)

**Official Symbol** [SIRPA](#)

**Synonyms** SIRPA; signal-regulatory protein alpha; protein tyrosine phosphatase, non receptor type substrate 1 , PTPNS1; tyrosine-protein phosphatase non-receptor type substrate 1; BIT; CD172a; MFR; MYD 1; P84; SHPS 1; SHPS1; SIRP; SIRP ALPHA 1; SIRPalpha; SIRPalpha2; myd-1 antigen; inhibitory receptor SHPS-1; macrophage fusion receptor; CD172 antigen-like family member A; tyrosine phosphatase SHP substrate 1; brain-immunoglobulin-like molecule with tyrosine-based activation motifs; MYD-1; CD172A; PTPNS1;


**Gene ID** [140885](#)

**mRNA Refseq** [NM\\_001040022](#)

**Protein Refseq** [NP\\_001035111](#)

**MIM** [602461](#)

**UniProt ID** [P78324](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA