

Recombinant Human SIRT2 293 Cell Lysate

Cat. No. SIRT2-1834HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for sirtuin (silent mating type information regulation 2 homolog) 2 (<i>S. cerevisiae</i>) (SIRT2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	SIRT2 sirtuin 2 [Homo sapiens]
Official Symbol	SIRT2
Synonyms	SIRT2; sirtuin 2; SIR2L, sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae) , sirtuin (silent mating type information regulation 2, S.cerevisiae, homolog) 2; NAD-dependent deacetylase sirtuin-2; sirtuin type 2; SIR2-like protein 2; sir2-related protein type 2; silent information regulator 2; SIR2; SIR2L; SIR2L2; FLJ35621; FLJ37491;
Gene ID	22933
mRNA Refseq	NM_030593
Protein Refseq	NP_085096
MIM	604480
UniProt ID	Q8IXJ6
Chromosome Location	19q13
Pathway	Signaling events mediated by HDAC Class I, organism-specific biosystem; Signaling events mediated by HDAC Class III, organism-specific biosystem;
Function	NOT NAD+ ADP-ribosyltransferase activity; NAD+ binding; NAD-dependent histone

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



deacetylase activity; histone acetyltransferase binding; histone deacetylase binding; hydrolase activity; metal ion binding; protein binding; protein deacetylase activity; transcription factor binding; tubulin deacetylase activity; ubiquitin binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA