

Recombinant Human SIRT2

Cat. No. SIRT2-30564TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Full-length recombinant Human SIRT2 with an N-terminal tag. MWt 64kDa.
Species	Human
Source	E.coli
ProteinLength	389 amino acids
Description	<p>This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative splicing of this gene.</p>
Molecular Weight	64.000kDa inclusive of tags
Tissue specificity	<p>Widely expressed. Highly expressed in heart, brain and skeletal muscle, while it is weakly expressed in placenta and lung. Down-regulated in many gliomas suggesting that it may act as a tumor suppressor gene in human gliomas possibly through the regulatio</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Liquid
Storage buffer	Preservative: None Constituents: 25% Glycerol, 50mM Tris HCl, 150mM Sodium chloride, 10mM Glutathione, 0.25mM DTT, 0.1mM EDTA, 0.1mM PMSF, pH 7.5
Storage	Shipped on dry ice. Upon delivery aliquot and store at -80oC. Avoid freeze / thaw cycles.
Sequence Similarities	Belongs to the sirtuin family. Contains 1 deacetylase sirtuin-type domain.

GENE INFORMATION

Gene Name	SIRT2 sirtuin 2 [Homo sapiens]
Official Symbol	SIRT2
Synonyms	SIRT2; sirtuin 2; SIR2L, sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae) , sirtuin (silent mating type information regulation 2, S.cerevisiae, homolog) 2; NAD-dependent deacetylase sirtuin-2;
Gene ID	22933
mRNA Refseq	NM_001193286
Protein Refseq	NP_001180215
MIM	604480
Uniprot ID	Q8IXJ6
Chromosome	19q13

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Signaling events mediated by HDAC Class I, organism-specific biosystem; Signaling events mediated by HDAC Class III, organism-specific biosystem;
Function	NOT NAD+ ADP-ribosyltransferase activity; NAD+ binding; NAD-dependent histone deacetylase activity; histone acetyltransferase binding; histone deacetylase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA