

Active Recombinant Human SIRT3 protein, His-tagged

Cat. No. SIRT3-10H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human SIRT3 protein(Q9NTG7)(126-399 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	126-399 aa
Form	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	Sirtuin 3 (SIRT3), the NAD-dependent deacetylase is a member of the mammalian sirtuin family of proteins. In human, sirtuins have a range of molecular functions and have emerged as important proteins in aging, stress resistance and metabolic regulation. It also can regulate epigenetic gene silencing and suppress recombination of rDNA in yeast. SIRT3 expression in white and brown adipose tissue. Besides, Isocitrate Dehydrogenase2, mitochondrial (IDH2) has been identified as an interact or of SIRT3, thus a binding ELISA assay was conducted to detect the interaction of recombinant human SIRT3 and recombinant human IDH2. Briefly, SIRT3 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to IDH2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-SIRT3 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 μl stop solution to the wells and read at 450nm immediately. The binding activity of SIRT3 and IDH2 was shown, and this effect was in a dose dependent manner.

Molecular Mass 31.7kDa

Purity > 90%

Storage Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name SIRT3 sirtuin 3 [Homo sapiens]

Official Symbol SIRT3

Synonyms SIRT3; sirtuin 3; sirtuin (silent mating type information regulation 2 homolog) 3 (S. cerevisiae) , sirtuin (silent mating type information regulation 2, S.cerevisiae, homolog) 3; NAD-dependent deacetylase sirtuin-3, mitochondrial; SIR2L3; sir2-like 3; sirtuin type 3; SIR2-like protein 3; silent mating type information regulation 2, S.cerevisiae, homolog 3; mitochondrial nicotinamide adenine dinucleotide-dependent deacetylase;

Gene ID 23410

mRNA Refseq NM_001017524

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_001017524


MIM 604481

UniProt ID Q9NTG7

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA