

## Recombinant Human SKP1 293 Cell Lysate

Cat. No. SKP1-1813HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for S-phase kinase-associated protein 1 (SKP1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** SKP1 S-phase kinase-associated protein 1 [ Homo sapiens ]

**Official Symbol** SKP1

**Synonyms** SKP1; S-phase kinase-associated protein 1; S phase kinase associated protein 1A (p19A) , SKP1A; EMC19; MGC34403; OCP II; OCP2; p19A; TCEB1L; SIII; OCP-2; p19skp1; organ of Corti protein 2; organ of Corti protein II; cyclin A/CDK2-associated p19; transcription elongation factor B; cyclin A/CDK2-associated protein p19; cyclin-A/CDK2-associated protein p19; RNA polymerase II elongation factor-like protein OCP2; SKP1A; OCP-II;

**Gene ID** 6500

**mRNA Refseq** NM\_006930

**Protein Refseq** NP\_008861

**MIM** 601434

**UniProt ID** P63208

**Chromosome Location** 5q31

**Pathway** APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Proteasome degradation, organism-specific biosystem; Canonical Wnt signaling pathway, organism-specific biosystem; Cell Cycle, organism-specific biosystem;

**Function**

protein binding; ubiquitin-protein ligase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA