

Recombinant Human SLC7A10 293 Cell Lysate

Cat. No. SLC7A10-1699HCL Lot. No. (See product label)

SPECIFICATION

| | |
|----------------------------|---|
| Species | Human |
| Source | HEK293 |
| Description | Antigen standard for solute carrier family 7, (neutral amino acid transporter, y+ system) member 10 (SLC7A10) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| Components | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| Size | 0.1 mg |
| Storage Instruction | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |
| Applications | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name SLC7A10 solute carrier family 7 (neutral amino acid transporter light chain, asc system), member 10 [Homo sapiens]

Official Symbol SLC7A10

Synonyms asc-1; HASC-1

Gene ID 56301

mRNA Refseq NM_019849.2

Protein Refseq NP_062823.1

MIM 607959

UniProt ID Q9NS82

Chromosome Location 19q13.1

Pathway Amino acid transport across the plasma membrane, organism-specific biosystem;Basigin interactions, organism-specific biosystem;Cell surface interactions at the vascular wall, organism-specific biosystem;Hemostasis, organism-specific biosystem;SLC-mediated transmembrane transport, organism-specific biosystem;Transmembrane transport of small molecules, organism-specific biosystem;Transport of inorganic cations/anions and amino acids/oligopeptides, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

L-serine transmembrane transporter activity;neutral amino acid transmembrane transporter activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA