

## Active Recombinant Human STAT1, His-tagged

Cat. No. STAT1-8473H Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Recombinant Human STAT1, fused with His tag, was expressed in Sf21 Cells.

**Species** Human

**Source** Sf21 Cells

#### Description

Signal transducer and activator of transcription (STAT) proteins are a family of latent cytoplasmic transcription factors involved in cytokine, hormone, and growth factor signal transduction. Seven members of the STAT family of transcription factors have been identified in mammalian cells: STAT1, STAT2, STAT3, STAT4, STAT5a, STAT5b, and STAT6. STAT proteins mediate broadly diverse biologic processes, including cell growth, differentiation, apoptosis, fetal development, transformation, inflammation, and immune response. Receptor-recruited STATs are phosphorylated on a single tyrosine residue in the carboxy terminal portion. The modified STATs are released from the cytoplasmic region of the receptor subunits to form homodimers or heterodimers through reciprocal interaction between the phosphotyrosine of one STAT and the SH2 domain of another. Following dimerization, STATs rapidly translocate to the nucleus and interact with specific regulatory elements to induce target gene transcription. Recently, STAT-1 has been implicated in modulating pro- and anti-apoptotic genes following several stress-induced responses. These effects are dependent on STAT-1 phosphorylation on serine-727 and require the C-terminal transactivation domain of STAT-1 to enhance its pro-apoptotic effect or inhibit its anti-apoptotic effects. The STAT-1 C-terminal domain has been demonstrated to be important for protein-protein interaction with other transcriptional activators. The

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reports that STAT-1-deficient mice develop spontaneous and chemically induced tumours more rapidly compared to wild-type mice and that STAT-1-deficient cells are more resistant to agents that induce apoptosis strongly support the argument that STAT-1 acts as a tumour suppressor.

**Form** 20 mM Tris-Cl, pH 7.9, 20% Glycerol, 100 mM KCl, 1 mM DTT and 0.2 mM EDTA

**Bio-activity** 1 unit equals 1 ng of purified protein. 1 unit is the amount sufficient for a gel mobility shift assay in a 20 µl reaction; 100 units are sufficient for a protein-protein interaction assay detected by an immuno-blot system.

**AA Sequence**

```

MSQWYELQQL DSKFLEQVHQ LYDDSPFMEI RQYLAQWLEK QDWEHAANDV
SFATIRFHDL LSQLDDQYSR FSLENNFLLQ HNIRKSKRNL QDNFQEDPIQ
MSMIIYSCLK EERKILENAQ RFNQAQSGNI QSTVMLDKQK ELDSKVRNVK
DKVMCIEHEI KSLEDLQDEY DFKCKTLQNR EHETNGVAKS DQKQEQLLLK
KMYLMLDNKR KEVVHKIIEI LNVTELTQNA LINDELVEWK RRQQSACIGG
PPNACLDQLQ NWFIVAESL QQVRQQLKKL EELEQKYTYE HDPITKNKQV
LWDRTFSLFQ QLIQSSFVVE RQPCMPHPQ RPLVLKTGVQ FTVKLRLLLVK
LQELNYNLKV KVLFDKDVNE RNTVKGFRKF NILGHTTKVM NMEESTNGSL
AAEFRHLQLK EQKNAGTRTN EGPLIVTEEL HSLSFETQLC QPGLVIDLET
TSLPVVVISN VSQLPSGWAS ILWYNMLVAE PRNLSFFLTP PCARWAQLSE
VLSWQFSSVT KRGLNVDQLN MLGEKLLGPN ASPDGLIPWT RFCKENINDK
NFPFWLWIES ILELIKKHLL PLWNDGCIMG FISKERERAL LKDQQPGTFL
LRFSESSREG AITFTWVERS QNGGEPDFHA VEPYTKKELS AVTFPDIIRN
YKVMAAENIP ENPLKYLYPN IDKDHAFGKY YSRPKEAPEP MELDGPKG TG
YIKTELISVS EVHPSRLQTT DNLLPMSPEE FDEVSRIVGS VEFDSMMNTV
    
```

**Purity** >95% as determined by SDS-PAGE

**Storage** -80 °C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Shipping** Dry Ice

## GENE INFORMATION

**Gene Name** [STAT1 signal transducer and activator of transcription 1, 91kDa \[ Homo sapiens \]](#)

**Official Symbol** STAT1

**Synonyms** STAT1; signal transducer and activator of transcription 1, 91kDa; signal transducer and activator of transcription 1, 91kD; signal transducer and activator of transcription 1-alpha/beta; ISGF 3; STAT91; transcription factor ISGF 3 components p91/p84; transcription factor ISGF-3 components p91/p84; signal transducer and activator of transcription-1; CANDF7; ISGF-3; DKFZp686B04100;

**Gene ID** [6772](#)

**mRNA Refseq** [NM\\_007315](#)

**Protein Refseq** [NP\\_009330](#)

**MIM** [600555](#)

**UniProt ID** [P42224](#)

**Chromosome Location** 2q32.2-q32.3

**Pathway** Adipogenesis, organism-specific biosystem; Antiviral mechanism by IFN-stimulated genes, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

pathway, conserved biosystem; Cytokine Signaling in Immune system, organism-specific biosystem;

**Function**

RNA polymerase II core promoter sequence-specific DNA binding; RNA polymerase II core promoter sequence-specific DNA binding transcription factor activity; calcium ion binding; double-stranded DNA binding; enzyme binding; non-membrane spanning protein tyrosine kinase activity; protein binding; protein homodimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; NOT sequence-specific DNA binding transcription factor activity; signal transducer activity; tumor necrosis factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA