

Recombinant Human STAT3 protein, His-tagged

Cat. No. STAT3-31H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human STAT3 protein(Ile576~Pro678), fused with N-terminal His tag, was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	576-678 a.a.
Form	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose.

Bio-activity	<p>Signal transducer and activator of transcription 3 (STAT3) is a member of the STAT protein family. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. STAT3 is phosphorylated by receptor-associated Janus kinases (JAK), form homo- or heterodimers, and translocate to the cell nucleus where they act as transcription activators. Besides, Epidermal Growth Factor Receptor 2 (EGFR2) has been identified as an interactor of STAT3, thus a binding ELISA assay was conducted to detect the interaction of recombinant human STAT3 and recombinant human EGFR2. Briefly, STAT3 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100µL were then transferred to EGFR2-coated microtiter wells and incubated for 2h at 37°C . Wells were washed with PBST and incubated for 1h with anti-STAT3 pAb, then aspirated and washed 3 times. After incubation with HRP</p>
---------------------	--

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of STAT3 and EGFR2 was shown in Figure, and this effect was in a dose dependent manner.

Molecular Mass 12.9kDa

Endotoxin <1.0EU per 1µg (determined by the LAL method).

Purity >95%

Storage Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name STAT3 signal transducer and activator of transcription 3 (acute-phase response factor) [Homo sapiens]

Official Symbol STAT3

Synonyms STAT3; signal transducer and activator of transcription 3 (acute-phase response factor); signal transducer and activator of transcription 3; APRF; DNA-binding protein APRF; acute-phase response factor; HIES; FLJ20882; MGC16063;

Gene ID 6774

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM_003150](#)

Protein Refseq [NP_003141](#)

MIM [102582](#)

UniProt ID [P40763](#)

SDS-PAGE




 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

WB



Activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA