

Recombinant Human STAT6 cell lysate

Cat. No. STAT6-481HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human STAT6 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

STAT6 signal transducer and activator of transcription 6, interleukin-4 induced [Homo sapiens]

Official Symbol

STAT6

Synonyms

STAT6; signal transducer and activator of transcription 6, interleukin-4 induced; signal transducer and activator of transcription 6; D12S1644; IL 4 STAT; STAT, interleukin4-induced; transcription factor IL-4 STAT; STAT6B; STAT6C; IL-4-STAT;

Gene ID

6778

mRNA Refseq

NM_001178078

Protein Refseq

NP_001171549

MIM

601512

UniProt ID

P42226

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	12q13
Pathway	Adipogenesis, organism-specific biosystem; Downstream signal transduction, organism-specific biosystem; IL-3 Signaling Pathway, organism-specific biosystem; IL-4 signaling Pathway, organism-specific biosystem; IL12-mediated signaling events, organism-specific biosystem; IL4-mediated signaling events, organism-specific biosystem; Jak-STAT signaling pathway, organism-specific biosystem;
Function	calcium ion binding; identical protein binding; protein binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA