

Recombinant Human Stanniocalcin 1, Flag-tagged

Cat. No. STC1-132H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human Stanniocalcin-1 produced in 293 cell line is a single, glycosylated, polypeptide chain containing 240 amino acids and having a total molecular mass of 25.9 kDa. The Stanniocalcin contains two extra residues which were used as a spacer and 8 residues form the C-Terminal Flag- tag. Stanniocalcin is purified by proprietary chromatographic techniques. The amino acid sequence of the recombinant human Stanniocalcin-1 is 100% homologous to the amino acid sequence AA 18-247 of the human mature Human Stanniocalcin-1.
Species	Human
Source	HEK293
Description	Stanniocalcin 1 (STC1) is the mammalian homologue of STC, which was originally identified as a calcium/phosphate-regulating hormone in bony fishes. In contrast, STC1 may play an autocrine and paracrine role with pleiotropic effects in mammals. It is expressed in a wide variety of tissues, but unexpectedly is not detected in the circulation under normal circumstances, which is possibly caused by its attaching to soluble and tethered forms of a high-affinity binding protein. STC-1 can affect calcium homeostasis, bone and muscle mass and structure, and angiogenesis through effects on osteoblasts, osteoclasts, myoblasts/myocytes, and endothelial cells in mouse model. Differential regulation of myocardial STC1 protein expression was reported in heart failure. In addition, STC1 may regulate calcium currents in cardiomyocytes and may contribute to the alterations in calcium homeostasis of the failing heart. STC1 was found to be a selective modulator of hepatocyte growth factor (HGF)-induced

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

endothelial migration and morphogenesis, an inhibitor of macrophage chemotaxis and chemokinesis, suppressor of progesterone and luteinization inhibitor. Together with STC-2, it may play important roles in the processes of implantation and decidualization in the rat. In terminally differentiated adipocytes, it may function as a "survival factor", which contributes to the maintenance of the integrity of mature adipose tissue. In context with its possible role in gestation, a Big STC, a three higher-molecular-mass variant has been described. STC1 was identified as one of hypoxia-responsive genes coupled to hypoxia-driven angiogenesis. Current research indicates that STC-1 might be a useful molecular marker to detect tumor cells in blood and bone marrow from patients with various types of malignancies.

Amino Acid Sequence

THEAEQNDSV SPRKSRVAAQ NSAEVVRCLN SALQVGCGAF ACLENSTCDT
 DGMVDICKSF LYSAKFDTQ GKAFVKESLK CIANGVTSKV FLAIRRCSTF
 QRMIAEVQEE CYSKLVNCSI AKRNPEAITE VVQLPNHFSN RYYNRLVRS
 LECDEDTVST IRDSLMEKIG PNMASLFHIL QTDHCAQTHP RADFNRRRTN
 EPQKLVLLR NLRGEEDSPS HIKRTSHESA ASDYKDDDDK

Physical Appearance

White lyophilized (freeze-dried) powder.

Form

Filtered (0.4 μm) and lyophilized in 0.5 mg/ml in 20mM Tris buffer, 20mM NaCl, pH 7.5.

Purity

Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Solubility


Add deionized water to a working concentration approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by appropriate sterile filter before using it in the cell culture.

Stability

Lyophilized STC-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Stanniocalcin should be stored at 4°C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

GENE INFORMATION

Gene Name	STC1 stanniocalcin 1 [Homo sapiens]
Synonyms	STC1; stanniocalcin 1; STC; Stanniocalcin-1; STC-1
Gene ID	6781
mRNA Refseq	NM_003155
Protein Refseq	NP_003146
MIM	601185
UniProt ID	P52823
Chromosome Location	8p22-p12
Function	hormone activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA