

Recombinant Human TAOK3 cell lysate

Cat. No. TAOK3-587HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human TAOK3 / MAP3K18 (aa 1 - 411) derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
ProteinLength	1-411 a.a.
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

TAOK3 TAO kinase 3 [[Homo sapiens](#)]

Official Symbol

TAOK3

Synonyms

TAOK3; TAO kinase 3; serine/threonine-protein kinase TAO3; DPK; JIK; MAP3K18; hKFC-A; STE20-like kinase; JNK/SAPK-inhibitory kinase; jun kinase-inhibitory kinase; kinase from chicken homolog A; CTCL-associated antigen HD-CL-09; dendritic cell-derived protein kinase; thousand and one amino acid protein 3; cutaneous T-cell lymphoma-associated antigen HD-CL-09; FLJ31808; DKFZp666H245;

Gene ID

[51347](#)

mRNA Refseq

[NM_016281](#)

Protein Refseq

[NP_057365](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



UniProt ID	Q9H2K8
Chromosome Location	12q
Pathway	MAPK signaling pathway, organism-specific biosystem; MAPK signaling pathway, conserved biosystem; p38 MAPK signaling pathway, organism-specific biosystem;
Function	ATP binding; nucleotide binding; protein kinase inhibitor activity; protein serine/threonine kinase activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA