

Recombinant Human TFDP1 293 Cell Lysate

Cat. No. TFDP1-1132HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for transcription factor Dp-1 (TFDP1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [TFDP1 transcription factor Dp-1 \[Homo sapiens \]](#)

Official Symbol [TFDP1](#)

Synonyms [TFDP1](#); transcription factor Dp-1; Dp 1; DP1; DRTF1; DRTF1-polypeptide 1; E2F dimerization partner 1; E2F-related transcription factor; Dp-1;

Gene ID [7027](#)

mRNA Refseq [NM_007111](#)

Protein Refseq [NP_009042](#)

MIM [189902](#)

UniProt ID [Q14186](#)

Chromosome Location [13q34](#)

Pathway [Activation of BH3-only proteins, organism-specific biosystem](#); [Activation of NOXA and translocation to mitochondria, organism-specific biosystem](#); [Activation of PUMA and translocation to mitochondria, organism-specific biosystem](#); [Apoptosis, organism-specific biosystem](#); [Cell Cycle, organism-specific biosystem](#); [Cell Cycle, Mitotic, organism-specific biosystem](#); [Cell cycle, organism-specific biosystem](#);

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA