

## Active Recombinant Human TFF2 protein, Tag Free

Cat. No. TFF2-30657TH Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human TFF2 protein was expressed in Escherichia coli.
<b>Species</b>	Human
<b>Source</b>	E.coli
<b>ProteinLength</b>	24-129 a.a.
<b>Description</b>	Trefoil factor 2 encoded by the TFF2 gene in humans, belongs to the trefoil factor family that consists of three members named TFF1, TFF2 and TFF3. They are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfides. The TFFs are stable secretory proteins expressed highly in the gastrointestinal tract (gastric mucosa). TFF2 can inhibit gastrointestinal motility and gastric acid secretion. Additionally, it functions as a structural component of gastric mucus, possibly by stabilizing glycoproteins in the mucus gel through interactions with carbohydrate side chains.
<b>Form</b>	Lyophilized from a 0.2µm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl.
<b>Bio-activity</b>	Fully biologically active when compared to standard. The ED50 as determined by a chemotaxis bioassay using human MCF-7 cells is less than 10 ng/ml, corresponding to a specific activity of > 1.0 × 10 <sup>5</sup> IU/mg.
<b>Molecular Mass</b>	Approximately 12.0 kDa, a single non-glycosylated polypeptide chain containing 106

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

amino acids, which includes a 40-amino acid trefoil motif containing three conserved intramolecular disulfide bonds.

**AA Sequence**

EKSPQCQCSRLSPHNRTNCGFPGITSDQCDFDNGCCFDSSVTGVPWCFFHPLPKQES  
DQCVMEVSDRRNCGYPGISPEECASRKCCFSNFIFEVWCFFPKSVEDCHY

**Endotoxin**

Less than 1 EU/g of rHuTFF2 as determined by LAL method.

**Purity**

>97% by SDS-PAGE and HPLC analysis.

**Storage**

Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

## GENE INFORMATION

**Gene Name**

TFF2

**Official Symbol**

TFF2

**Synonyms**

TFF2; trefoil factor 2; SML1, spasmolytic protein 1; spasmolysin; spasmolytic protein 1; spasmolytic polypeptide; SP; SML1;

**Gene ID**

7032

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>mRNA Refseq</b>	NM_005423
<b>Protein Refseq</b>	NP_005414
<b>MIM</b>	182590
<b>UniProt ID</b>	Q03403

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA