

Recombinant Mouse TFRC cell lysate

Cat. No. TFRC-1813MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse TFRC / CD71 / Transferrin Receptor derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Tfrc transferrin receptor [<i>Mus musculus</i>]
Official Symbol	TFRC
Synonyms	TFRC; transferrin receptor; transferrin receptor protein 1; mammary tumor virus receptor 1; TR; TFR; p90; CD71; TFR1; Trfr; Mtvr1; Mtvr-1; AI195355; AI426448; AU015758; 2610028K12Rik; E430033M20Rik;
Gene ID	22042
mRNA Refseq	NM_011638
Protein Refseq	NP_035768
Pathway	Clathrin derived vesicle budding, organism-specific biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved biosystem; Golgi Associated Vesicle Biogenesis, organism-specific biosystem; Hematopoietic cell lineage,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



organism-specific biosystem; Hematopoietic cell lineage, conserved biosystem;
Membrane Trafficking, organism-specific biosystem;

Function

Hsp70 protein binding; chaperone binding; iron ion transmembrane transporter
activity; peptidase activity; receptor activity; transferrin receptor activity; transferrin
receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA