

Recombinant Human TGFB11 Protein, His-tagged

Cat. No. TGFB111-652H Lot. No. (See product label)

SPECIFICATION

| | |
|-------------------------|---|
| Product Overview | Recombinant Human TGFB11(Phe252~Arg390+RCARTTSTHDAARCAPRVASL) fused with His tag at N-terminal was expressed in E. coli. |
| Species | Human |
| Source | E.coli |
| ProteinLength | Phe252~Arg390+RCARTTSTHDAARCAPRVASL |
| Description | This gene encodes a coactivator of the androgen receptor, a transcription factor which is activated by androgen and has a key role in male sexual differentiation. The encoded protein is thought to regulate androgen receptor activity and may have a role to play in the treatment of prostate cancer. Multiple transcript variants encoding different isoforms have been found for this gene. |
| Form | 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| Molecular Mass | 20.79kDa |
| Endotoxin | <1.0EU per 1g (determined by the LAL method) |
| Purity | > 92% |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

If bio-activity of the protein is needed, please check active protein.

Stability

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 centigrade for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Storage

Avoid repeated freeze/thaw cycles. Store at 2-8 centigrade for one month. Aliquot and store at -80 centigrade for 12 months.

Reconstitution

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name

TGFB111 transforming growth factor beta 1 induced transcript 1 [Homo sapiens]

Official Symbol

TGFB111

Synonyms

TGFB111; transforming growth factor beta 1 induced transcript 1; transforming growth factor beta-1-induced transcript 1 protein; ARA55; Hic 5; HIC 5; TSC 5; androgen receptor coactivator ARA55; hydrogen peroxide-inducible clone 5 protein; androgen receptor coactivator 55 kDa protein; androgen receptor-associated protein of 55 kDa; HIC5; HIC-5; TSC-5;

Gene ID

7041

mRNA Refseq

NM_001042454

Protein Refseq

NP_001035919


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM 602353

UniProt ID O43294



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA