

Recombinant human TGFB2, Active

Cat. No. TGFB2-1566H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human TGF- β 2 is a 27.08 kDa protein composed of two identical 118 amino acid polypeptide chains linked by a single disulphide bond.
Species	Human
Source	Nicotiana Benthamiana
Description	Recombinant human TGF- β 2 is a 27.08 kDa protein composed of two identical 118 amino acid peptide chains linked by a single disulphide bond. Transforming growth factor- β is a family of five related cytokines that have been shown on a wide variety of normal and neoplastic cells, indicating the importance of these homo-dimer proteins as multi-functional regulators of cellular activity. The three mammalian isoforms of TGF- β (TGF- β 1, TGF- β 2 and TGF- β 3) signal through the same receptor and elicit similar biological responses. They are involved in physiological processes as embryogenesis, tissue remodelling and wound healing.
Form	Recombinant human TGF-b2 is lyophilized from a Tris HCl 0.05M buffer at pH 7.4.
Molecular Mass	27.08 kDa
AA Sequence	HHHHHHALDAAYCFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAGACP YLWSSDTQHSRVLSLYNTIN PEASASPCCVSQDLEPLTILYYIGKTPKIEQLSNMIVK SCKCS
Endotoxin	< 0.04=""

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	eu/μg="" protein="" (lal="">
Purity	>97% by SDS-PAGE gel
Storage	This lyophilized preparation is stable at 2-8o C for short term, long storage it should be kept at -20oC. Reconstituted protein should be stored in working aliquots at -20°C. Repeated freezing and thawing is not recommended.
Reconstitution	Centrifuge vial before opening. Lyophilized protein should be reconstituted in water to a concentration of 25ng/ul. At higher concentration the solubility may be reduced and multimers generated.

GENE INFORMATION

Gene Name	TGFB2 transforming growth factor, beta 2 [Homo sapiens]
Official Symbol	TGFB2
Synonyms	TGFB2; transforming growth factor, beta 2; transforming growth factor beta-2; G-TSF; cetermin; polyergin; BSC-1 cell growth inhibitor; glioblastoma-derived T-cell suppressor factor; TGF-beta2; MGC116892;
Gene ID	7042
mRNA Refseq	NM_001135599
Protein Refseq	NP_001129071
MIM	190220
UniProt ID	P61812

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	1q41
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Cell cycle, organism-specific biosystem; Cell cycle, conserved biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem;
Function	beta-amyloid binding; cytokine activity; growth factor activity; protein binding; contributes_to protein binding; protein heterodimerization activity; protein homodimerization activity; receptor binding; receptor signaling protein serine/threonine kinase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA