

Recombinant Active Human TGFB2 Protein, His-tagged(C-ter)

Cat. No. TGFB2-302H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Active Human TGFB2 Protein with His tag (C-ter) was expressed in E. coli.
Species	Human
Source	E.coli
Description	<p>This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types by transducing their signal through combinations of transmembrane type I and type II receptors (TGFB1 and TGFB2) and their downstream effectors, the SMAD proteins. Disruption of the TGFB/SMAD pathway has been implicated in a variety of human cancers. The encoded protein is secreted and has suppressive effects of interleukin-2 dependent T-cell growth. Translocation t(1;7)(q41;p21) between this gene and HDAC9 is associated with Peters' anomaly, a congenital defect of the anterior chamber of the eye. The knockout mice lacking this gene show perinatal mortality and a wide range of developmental, including cardiac, defects. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2010]</p>
Form	Powder
Bio-activity	Measure by its ability to inhibit IL-4-induce proliferation in HT-2 cells. The ED50 for this effect is < 0.2 ng/mL. The specific activity of recombinant human TGF beta 2 is >

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	5 x 10 ⁶ IU/mg.
AA Sequence	MALDAAYCFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAGACPYLWSS DTQHRSRVLSLYNTINPEASASPCCVSDLEPLTILYIYGKTPKIEQLSNMIVKSKCS
Endotoxin	Endotoxin level is less than 0.01 EU/μg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	20 mM Sodium citrate (pH 3.5) and 0.2 M NaCl.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	TGFB2 transforming growth factor, beta 2 [Homo sapiens]
Official Symbol	TGFB2
Synonyms	TGFB2; transforming growth factor, beta 2; transforming growth factor beta-2; G-TSF;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cetermin; polyergin; BSC-1 cell growth inhibitor; glioblastoma-derived T-cell suppressor factor; TGF-beta2; MGC116892;

Gene ID 7042


mRNA Refseq NM_001135599

Protein Refseq NP_001129071

MIM 190220

UniProt ID P61812

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA