

Recombinant Human TGFBR1 cell lysate

Cat. No. TGFBR1-2482HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human TGFBR1 / ALK-5 / SKR4 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

TGFBR1 transforming growth factor, beta receptor 1 [Homo sapiens]

Official Symbol

TGFBR1

Synonyms

TGFBR1; transforming growth factor, beta receptor 1; transforming growth factor, beta receptor I (activin A receptor type II like kinase, 53kD); TGF-beta receptor type-1; activin A receptor type II like kinase; 53kDa; ACVRLK4; ALK 5; tbetaR-I; TGF-beta receptor type I; TGF-beta type I receptor; activin receptor-like kinase 5; transforming growth factor beta receptor I; serine/threonine-protein kinase receptor R4; activin A receptor type II-like kinase, 53kD; activin A receptor type II-like kinase, 53kDa; transforming growth factor-beta receptor type I; activin A receptor type II-like protein kinase of 53kD; transforming growth factor, beta receptor I (activin A receptor type II-like kinase, 53kD); AAT5; ALK5; MSSE; SKR4; ALK-5; LDS1A; LDS2A; TGFR-1;

Gene ID

7046

mRNA Refseq

NM_001130916

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_001124388
MIM	190181
UniProt ID	P36897
Chromosome Location	9q22
Pathway	ALK1 signaling events, organism-specific biosystem; Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chronic myeloid leukemia, organism-specific biosystem; Chronic myeloid leukemia, conserved biosystem;
Function	ATP binding; I-SMAD binding; SMAD binding; SMAD binding; contributes_to growth factor binding; metal ion binding; nucleotide binding; protein binding; protein heterodimerization activity; protein serine/threonine kinase activity; receptor activity; transforming growth factor beta binding; contributes_to transforming growth factor beta binding; transforming growth factor beta binding; transforming growth factor beta receptor activity, type I; transforming growth factor beta-activated receptor activity; transforming growth factor beta-activated receptor activity; transforming growth factor beta-activated receptor activity; contributes_to transforming growth factor beta-activated receptor activity; transmembrane receptor protein serine/threonine kinase activity; type II transforming growth factor beta receptor binding; type II transforming growth factor beta receptor binding; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA