

Recombinant Cynomolgus monkey TGFBR2 Protein, Fc&avi tagged, Biotinylated

Cat. No. TGFBR2-001CB **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated recombinant Cynomolgus monkey TGFBR2 Protein (24-159 aa) with Fc&avi tag was expressed in HEK293.
Species	Cynomolgus monkey
Source	HEK293
ProteinLength	24-159 aa
Conjugation/Label	Biotin
Description	<p>Transmembrane serine/threonine kinase forming with the TGF-beta type I serine/threonine kinase receptor, TGFBR1, the non-promiscuous receptor for the TGF-beta cytokines TGFB1, TGFB2 and TGFB3. Transduces the TGFB1, TGFB2 and TGFB3 signal from the cell surface to the cytoplasm and is thus regulating a plethora of physiological and pathological processes including cell cycle arrest in epithelial and hematopoietic cells, control of mesenchymal cell proliferation and differentiation, wound healing, extracellular matrix production, immunosuppression and carcinogenesis. The formation of the receptor complex composed of 2 TGFBR1 and 2 TGFBR2 molecules symmetrically bound to the cytokine dimer results in the phosphorylation and the activation of TGFRB1 by the constitutively active TGFBR2. Activated TGFBR1 phosphorylates SMAD2 which dissociates from the receptor and interacts with SMAD4. The SMAD2-SMAD4 complex is subsequently translocated to</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the nucleus where it modulates the transcription of the TGF-beta-regulated genes. This constitutes the canonical SMAD-dependent TGF-beta signaling cascade. Also involved in non-canonical, SMAD-independent TGF-beta signaling pathways.

Molecular Mass 44 kDa

AASequence
 IPPHVQKSVNNDMMVTDNNGAVKFPQLCKFCDVRFSTCDNQKSCLSNCSITSICEK
 PQEVCVAVWRKNDENITLETVCHDPKLPYHDFILEDAAAPKCIMKEKKKPGETFFMC
 SCSSDECNDNIIFSEEYNTSNPDIEGRMDPKSCDKTHTCPPCPAPPELLGGPSVFLFP
 PKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
 RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDE
 LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDK
 SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKGLNDIFEAQKIEWHE

Endotoxin <1 EU/μg by LAL

Purity > 90% by SDS-PAGE

Storage Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Concentration 2.05 mg/mL by BCA

Storage Buffer Sterile PBS, pH 7.4

Labeling efficiency 0.71 By HABA

GENE INFORMATION


Gene Name [TGFB2 transforming growth factor beta receptor 2 \[Macaca mulatta \(Rhesus monkey\) \]](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	TGFBR2
Synonyms	TGFBR2; transforming growth factor beta receptor 2; TGF-beta receptor type-2; transforming growth factor, beta receptor II (70/80kDa); EC 2.7.11.30
Gene ID	703088
mRNA Refseq	NM_001261151
Protein Refseq	NP_001248080
UniProt ID	H9YULO

SDS-PAGE and WB


Reducing 4%-20% SDS-PAGE (CBB stained) and WB (Goat Anti-Human IgG/HRP antibody) analysis profiles of purified TGFBR2. 1. TGFBR2: 1 µg 2. BSA: 1 µg

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA