

Active Recombinant Cynomolgus Monkey TIMD4 protein, Fc-tagged

Cat. No. TIMD4-198C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Cynomolgus Monkey TIMD4(Met22-Pro142) fused with Fc region of Human IgG1 at C-terminal was expressed in HEK293.
Species	Cynomolgus
Source	HEK293
ProteinLength	22-142 a.a.
Description	<p>TIM-4 (T cell; immunoglobulin; mucin-4), also known as SMUCKLER, is an approximately 60 kDa member of the TIM family of immune regulating proteins. TIMs are type I transmembrane proteins with one Ig-like V domain and one Ser/Thr-rich mucin domain. Sequence alignment with the human TIM-4 shows that mature cynomolgus monkey TIM-4 has a predicted 280 aa extracellular domain (ECD), a 21 aa transmembrane segment and a 43 aa cytoplasmic tail. Within the ECD, cynomolgus monkey TIM-4 shares 87 % and 48 % aa sequence identity with human and mouse TIM-4, respectively. Structurally, TIM-4 is distinguished from other TIMs by the presence of an RGD motif in its Ig domain and the lack of a site for tyrosine phosphorylation in its cytoplasmic tail. The mucin domain in TIM-4 is larger than in TIM-1 or TIM-3. TIM-4 is expressed by macrophages and mature dendritic cells but not by lymphocytes. TIM-4 binds specifically to TIM-1 which is also the cellular receptor for the hepatitis A virus, and has been implicated in the development of asthma. Among hematopoietic</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cells, TIM-1 is expressed on activated B and T cells, preferentially in the Th2 subset of CD4+ T cells. The interaction of TIM-4 with TIM-1 induces co-stimulatory and hyperproliferative signals in T cells.

Predicted N Terminal Glu25

Form Lyophilized from a 0.2 µm filtered solution in NaH₂PO₄, NaCl and EDTA.

Bio-activity Measured by its ability to inhibit anti-CD3-induced proliferation of stimulated human T cells. The ED50 for this effect is 0.2-1.2 µg/mL.

Molecular Mass Predicted Molecular Mass: 57 kDa
SDS-PAGE: 95-112 kDa, reducing conditions

Endotoxin <0.1 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
12 months from date of receipt, -20 to -70 centigrade as supplied.
1 month, 2 to 8 centigrade under sterile conditions after reconstitution.
3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

Reconstitution Reconstitute at 500 µg/mL in PBS.

GENE INFORMATION

Gene Name TIMD4 T-cell immunoglobulin and mucin domain containing 4 [*Macaca fascicularis*]

Official Symbol TIMD4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Synonyms

TIMD4; T-cell immunoglobulin and mucin domain containing 4

Gene ID

102140672

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA