

## Active Recombinant Human TIMD4 protein, Fc-tagged

Cat. No. TIMD4-244H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human TIMD4(Glu25-Gln314) fused Fc region of Human IgG1 was expressed in HEK293.
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	25-314 a.a.
<b>Description</b>	<p>TIM-4 (T cell; immunoglobulin; mucin-4), also known as SMUCKLER, is a 60 kDa member of the TIM family of immune regulating proteins. TIMs are type I transmembrane proteins with one Ig-like V domain and one Ser/Thr-rich mucin domain. Mature human TIM-4 consists of a 290 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 43 aa cytoplasmic tail. Structurally, TIM-4 is distinguished from other TIMs by the presence of an RGD motif in its Ig domain and the lack of a site for tyrosine phosphorylation in its cytoplasmic tail. Within the ECD human TIM-4 shares 45% aa sequence identity with mouse and rat TIM-4. TIM-4 is expressed by macrophages and mature dendritic cells but not by lymphocytes. It binds specifically to TIM-1 which is also the cellular receptor for hepatitis A virus and has been implicated in the development of asthma. Among hematopoietic cells, TIM-1 is expressed on activated B and T cells, preferentially in the Th2 subset of CD4+ T cells. The interaction of TIM-4 with TIM-1 induces co-stimulatory and hyperproliferative signals in T cells. TIM-4 also binds to phosphatidylserine (PS) on the surface of apoptotic cells. It enhances the clearance of apoptotic cells including T</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cells in the contraction phase of immune responses, apoptotic macrophages that accumulate in atherosclerosis, and dying neurons in the brain. TIM-4 is upregulated on some tumor cells and tumor-associated myeloid cells where it binds Integrin alpha V beta 3, contributes to tumor progression, and inhibits tumor-specific cellular immunity.

**Predicted N Terminal**    Glu25

**Form**    Lyophilized from a 0.2 µm filtered solution in PBS.

**Bio-activity**    Measured by its ability to inhibit anti-CD3-induced proliferation of stimulated human T cells. The ED50 for this effect is 0.15-0.75 µg/mL.

**Molecular Mass**    Predicted Molecular Mass: 58 kDa  
SDS-PAGE: 95-113 kDa, reducing conditions

**Endotoxin**    <0.1 EU per 1 µg of the protein by the LAL method.

**Purity**    >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

**Storage**    Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  
12 months from date of receipt, -20 to -70 centigrade as supplied.  
1 month, 2 to 8 centigrade under sterile conditions after reconstitution.  
3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

**Reconstitution**    Reconstitute at 500 µg/mL in PBS.

## GENE INFORMATION

**Gene Name**    [TIMD4 T-cell immunoglobulin and mucin domain containing 4 \[ Homo sapiens \]](#)

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Official Symbol</b>	TIMD4
<b>Synonyms</b>	TIMD4; T-cell immunoglobulin and mucin domain containing 4; T-cell immunoglobulin and mucin domain-containing protein 4; TIM-4; TIMD-4; T-cell membrane protein 4; TIM4; SMUCKLER; FLJ27515;
<b>Gene ID</b>	91937
<b>mRNA Refseq</b>	NM_001146726
<b>Protein Refseq</b>	NP_001140198
<b>MIM</b>	610096
<b>UniProt ID</b>	Q96H15

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA