

Recombinant Human TIMP1 cell lysate

Cat. No. TIMP1-2678HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human TIMP-1 / TIMP1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[TIMP1](#) [TIMP metalloproteinase inhibitor 1 \[Homo sapiens \]](#)

Official Symbol

TIMP1

Synonyms

TIMP1; TIMP metalloproteinase inhibitor 1; CLGI, TIMP, tissue inhibitor of metalloproteinase 1 (erythroid potentiating activity, collagenase inhibitor); metalloproteinase inhibitor 1; EPO; TIMP-1; collagenase inhibitor; erythroid potentiating activity; erythroid-potentiating activity; fibroblast collagenase inhibitor; tissue inhibitor of metalloproteinases 1; EPA; HCl; CLGI; TIMP; FLJ90373;

Gene ID

[7076](#)

mRNA Refseq

[NM_003254](#)

Protein Refseq

[NP_003245](#)

MIM

[305370](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P01033
Chromosome Location	Xp11.3-p11.23
Pathway	Activation of Matrix Metalloproteinases, organism-specific biosystem; Degradation of the extracellular matrix, organism-specific biosystem; Extracellular matrix organization, organism-specific biosystem; Hemostasis, organism-specific biosystem; IL6-mediated signaling events, organism-specific biosystem; Matrix Metalloproteinases, organism-specific biosystem; Platelet activation, signaling and aggregation, organism-specific biosystem;
Function	enzyme inhibitor activity; metal ion binding; metalloendopeptidase inhibitor activity; metalloendopeptidase inhibitor activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA