

Active Recombinant Human TIMP2 protein

Cat. No. TIMP2-29289TH Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Human TIMP2 protein was expressed in Escherichia coli.

Species Human


Source E.coli

ProteinLength 27-220 aa


Description This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix.

Form Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, with 3 % trehalose.

Bio-activity Fully biologically active when compared to standard. The biological activity as determined by its ability to inhibit human MMP-2 cleavage of a fluorogenic peptide substrate Mca-PLGL-Dpa-AR-NH₂.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 21.8 kDa, a single non-glycosylated polypeptide chain containing 194 amino acids.
AA Sequence	CSCSPVHPQQAFCNADVIRAKAVSEKEVDSGNDIYGNPIKRIQYEIKQIKMFKGPEK DIEFIYTAPSSAVCGVSLDVGGKKEYLIAGKAEGDGKMHITLCDFIVPWDTLSTTQKK SLNHRYQMGCECKITRCPMIPCYISSPDECLWMDWVTEKNINGHQAKFFACIKRSD GSCAWYRGAAPPKQEFLDIEDP
Endotoxin	Less than 1 EU/g of rHuTIMP-2 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	TIMP2
Official Symbol	TIMP2
Synonyms	TIMP2; TIMP metalloproteinase inhibitor 2; tissue inhibitor of metalloproteinase 2; metalloproteinase inhibitor 2; CSC 21K; TIMP-2; tissue inhibitor of metalloproteinases

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



2; CSC-21K;

Gene ID

7077

mRNA Refseq

NM_003255

Protein Refseq


NP_003246

MIM


188825

UniProt ID

P16035

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA