

Active Recombinant Human TIMP2 protein, His-TrxA-tagged

Cat. No. TIMP2-3485H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human TIMP2 protein(P16035)(27~220 aa), fused with His tag and TrxA tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	27-220 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). There are four members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4. TIMP-2 is a non N-glycosylated protein with a molecular mass of 22 kDa produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes. TIMP-2 also has erythroid potentiating and cell growth promoting activities. The activity of recombinant human TIMP2 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH ₂ in the assay buffer 50 mM Tris, 10 mM CaCl ₂ , 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM A PMA at 37°C for 1 hour and rhTIMP2 (MW: 51.75 KD) was diluted to different concentrations with the assay buffer. Mix 8 μof rhTIMP2 curve dilutions, 12.8 μof activated rh

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MMP-2, and 59.2 µof assay buffer, including a control containing assay buffer and the diluted rhMMP-2 and incubate the reactions for 2 hours at 37°C. Loading 50 µof the incubated mixtures which were diluted five-fold in assay buffer into empty wells of a plate, and start the reaction by adding 50 µof 20 M substrate. Include a substrate blank containing 50 µof assay buffer and 50 µof 20 M substrate. Then read at excitation and emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that recombinant human TIMP2 significantly decreased rhMMP2 activity. The inhibition IC50 was <20 nM.

Molecular Mass	42.2kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 80% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	TIMP2 TIMP metalloproteinase inhibitor 2 [Homo sapiens]
Official Symbol	TIMP2
Synonyms	TIMP2; TIMP metalloproteinase inhibitor 2; tissue inhibitor of metalloproteinase 2; metalloproteinase inhibitor 2; CSC 21K; TIMP-2; tissue inhibitor of metalloproteinases 2; CSC-21K;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 7077


mRNA Refseq NM_003255

Protein Refseq NP_003246


MIM 188825

UniProt ID P16035

SDS-PAGE



Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA