

## Active Recombinant Bovine TIMP3 protein, His-tagged

Cat. No. TIMP3-1271C Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Active Recombinant Bovine TIMP3 protein(P79121)(24~211 aa), fused with N-terminal His tag, was expressed in E.coli.
<b>Species</b>	Bovine
<b>Source</b>	E.coli
<b>ProteinLength</b>	24-211 aa
<b>Form</b>	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
<b>Bio-activity</b>	Tissue Inhibitors Of Metalloproteinase 3 (TIMP3) is a protein belongs to the tissue inhibitor of metalloproteinases family. They are inhibitors of the matrix metalloproteinases. TIMP-3 is the only member of the TIMP family which is found exclusively in the extracellular matrix (ECM). It is regulated in a cell cycle-dependent fashion in certain cell types and may serve as a marker for terminal differentiation. The activity of recombinant bovine TIMP3 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH <sub>2</sub> in the assay buffer 50 mM Tris, 10 mM CaCl <sub>2</sub> , 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM APMA at 37°C for 1 hour and rbTIMP3 (MW: 23.25 KD) was diluted to different concentrations with the assay buffer. Mix 8 μof rbTIMP3 curve dilutions, 12.8 μof activated rhMMP-2, and 59.2 μof assay buffer, including a control containing assay buffer and the diluted rhMMP-2 and incubate the reactions

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

for 2 hours at 37°C. Loading 50 µof the incubated mixtures which were diluted five-fold in assay buffer into empty wells of a plate, and start the reaction by adding 50 µof 200 µM substrate. Include a substrate blank containing 50 µof assay buffer and 50 µof 200 µM substrate. Then read at excitation and emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that recombinant bovine TIMP3 significantly decreased rhMMP2 activity. The inhibition IC50 was <15 nM.

**Molecular Mass** 25.4kDa


**Endotoxin** <1.0EU per 1µg (determined by the LAL method).

**Purity** > 95% as determined by SDS-PAGE.

**Storage** Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

**Reconstitution** Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

**SDS-PAGE**




 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Activity**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA