

## Active Recombinant Bovine TIMP4 protein, His-tagged

Cat. No. TIMP4-3584B Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Active Recombinant Bovine TIMP4 protein(O97563)(50~224 aa), fused with N-terminal His tag, was expressed in E.coli.
<b>Species</b>	Bovine
<b>Source</b>	E.coli
<b>ProteinLength</b>	50-224 aa
<b>Form</b>	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
<b>Bio-activity</b>	Tissue Inhibitors of Metalloproteinase 4 (TIMP4) is an enzyme that in humans is encoded by the TIMP4 gene. This gene belongs to the tissue inhibitor of metalloproteinases gene family. The proteins encoded by this gene family are inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. The secreted, netrin domain-containing protein encoded by this gene is involved in regulation of platelet aggregation and recruitment and may play role in hormonal regulation and endometrial tissue remodeling. The activity of recombinant bovine TIMP4 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH <sub>2</sub> in the assay buffer 50 mM Tris, 10 mM CaCl <sub>2</sub> , 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM APMA at 37°C for 1 hour and rbTIMP4 (MW: 24.05 KD) was diluted to different concentrations with the assay buffer. Mix 8 μof rbTIMP4 curve di

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

lutions, 12.8  $\mu$ of activated rhMMP-2, and 59.2  $\mu$ of assay buffer, including a control containing assay buffer and the diluted rhMMP-2 and incubate the reactions for 2 hours at 37°C. Loading 50  $\mu$ of the incubated mixtures which were diluted five-fold in assay buffer into empty wells of a plate, and start the reaction by adding 50  $\mu$ of 20  $\mu$ M substrate. Include a substrate blank containing 50  $\mu$ of assay buffer and 50  $\mu$ of 20  $\mu$ M substrate. Then read at excitation and emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that recombinant bovine TIMP4 significantly decreased rhMMP2 activity. The inhibition IC50 was <27 nM.

**Molecular Mass** 24.1kDa

**Endotoxin** <1.0EU per 1 $\mu$ g (determined by the LAL method).

**Purity** > 95% as determined by SDS-PAGE.

**Storage** Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

**Reconstitution** Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA



SDS-PAGE



Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA