

Active Recombinant Rhesus monkey TNF protein, His-tagged

Cat. No. TNF-4218R Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Rhesus monkey TNF protein(P48094)(77~197 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Rhesus monkey
Source	E.coli
ProteinLength	77-197 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>Tumor necrosis factor (TNF, tumor necrosis factor alpha, TNFα, cachexin, orcachectin) is a cell signaling protein (cytokine) involved in systemic inflammationand is one of the cytokines that make up the acute phase reaction. Theprimaryrole of TNF is in the regulation of immune cells. TNF, being an endogenouspyrogen, is able to induce fever, apoptotic cell death, cachexia, inflammation and to inhibit tumorigenesis and viral replication and respond to sepsis viaIL1&IL6 producing cells. To test the effect of TNFα on cell apoptosis, A549 cells were seeded into 96-well plates at a density of 4,000 cells/well with 5% serum standard DMEM including various concentrations of recombinant simian TNFα. After incubated for 48h, cells were observed by inverted microscopeand cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10μof CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating</p>


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the plate for 1h at 37°C. Proliferation of A549 cells after incubation with TNF α for 72h observed by inverted microscope was shown. Cell viability was assessed by CCK-8(Cell Counting Kit-8) assay after incubation with recombinant simian TNF α for 72h. It was obvious that TNF α significantly inhibit cell viability of A549 cells. The ED50 is 6.9 μ g/mL.

Molecular Mass	20.1kDa
Endotoxin	<1.0EU per 1 μ g (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


SDS-PAGE


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA