

Recombinant Human TNFRSF10D

Cat. No. TNFRSF10D-1693H **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	Mammalian Cells
Description	TNF-related apoptosis-inducing ligand receptor 4 (TRAIL R4) is a decoy receptor that negatively regulate TRAIL-induced cytotoxicity by competing for ligand binding with TRAIL-R1 and TRAIL-R2.1 TRAIL R4 is expressed mainly on CD8+ and NK cells and induces apoptosis in many tumour cells, but not in normal cells. TRAIL-R4 contains partially truncated death domain and therefore is unable to induce apoptosis and serves as a negative regulator of apoptotic signaling by impairment death-inducing signaling complex (DISC) processing.
Molecular Weight	The predicted molecular weight of Recombinant Human TRAIL R4 is Mr 44.2 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 70-80 kDa.
State Of Matter	Lyophilized.
Purity	>95% by SDS Page and analyzed by silver stain.
Endotoxin	<1.0 EU/g as determined by the LAL method.
Storage And Stability	This lyophilized protein is stable for six to twelve months when stored desiccated at -20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C for one month or at -20°C to -70°C in a manual defrost freezer. Avoid

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Repeated Freeze Thaw Cycles. See Product Insert for exact lot specific storage instructions.

GENE INFORMATION

Gene Name

TNFRSF10D tumor necrosis factor receptor superfamily, member 10d, decoy with truncated death domain [Homo sapiens]

Synonyms

TNFRSF10D; tumor necrosis factor receptor superfamily, member 10d, decoy with truncated death domain; DCR2; CD264; TRUNDD; TRAILR4; tumor necrosis factor receptor superfamily, member 10d; TRAIL receptor 4; decoy receptor 2; decoy with truncated death domain; TNF receptor-related receptor for TRAIL; TRAIL receptor with a truncated death domain; DcR2; TRAIL-R4; CD264 antigen; TNF-related apoptosis-inducing ligand receptor 4

Gene ID

8793

mRNA Refseq

NM_003840

Protein Refseq

NP_003831

MIM

603614

UniProt ID

Q9UBN6

Chromosome Location

8p21

Pathway

Apoptosis; Cytokine-cytokine receptor interaction; Natural killer cell mediated cytotoxicity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

protein binding; transmembrane receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA