

# Active Recombinant Human Tumor Necrosis Factor Receptor Superfamily, Member 11b

**Cat. No.** TNFRSF11B-4141H    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	Recombinant OPG produced in yeast contains 412 amino acid residues, including 180 residues from mature OPG (a.a 22-201) and 232 residues from the Fc protein of human IgG1, and has a calculated molecular mass of 46.5 kDa. As a result of glycosylation, the recombinant Osteoprotegrin migrates as a 49 kDa protein in SDS-PAGE under reducing conditions. The OPG is purified by proprietary chromatographic techniques.
<b>Species</b>	Human
<b>Source</b>	Human
<b>ProteinLength</b>	22-201 a.a.
<b>Description</b>	Osteoprotegerin acts as decoy receptor for rankl and thereby neutralizes its function in osteoclastogenesis. OPG inhibits the activation of osteoclasts and promotes osteoclast apoptosis in vitro. Bone homeostasis seems to depend on the local rankl/opg ratio. Osteoprotegerin may also play a role in preventing arterial calcification. May act as decoy receptor for trail and protect against apoptosis. Trail binding blocks the inhibition of osteoclastogenesis.
<b>Form</b>	OPG was lyophilized from a 0.2µm filtered concentrated (0.5mg/ml) solution in PBS, pH= 7.4.
<b>Purity</b>	Greater than 90.0% as determined by(a) Analysis by RP-HPLC. (b) Analysis by SDS-

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	PAGE.
<b>Bio-activity</b>	Determined by its ability to neutralize the stimulation of U937 cells treated with 10ng/ml of soluble RANKL (sRANKL) corresponding to a Specific Activity of 100,000IU/mg.
<b>Physical Appearance</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Solubility</b>	It is recommended to reconstitute the lyophilized Osteoprotegerin in sterile 18M $\mu$ -cm H <sub>2</sub> O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
<b>Amino acid sequence</b>	AA Sequence: OPG 22-201: ETFPPKYLHYDEETSHQLLCDKCPPGTYLKQHCTAKW KTVCAPCPDHY, TDSWHTSDECLYCSPVCKELQYVKQECNRTHNRVCECKEGRY LEIEFCL, KHRSCPPGFGVVQAGTPERNVCKRCPDGGFFSNETSSKAPCRKHTNCS, VFGLLLTQKGNATHDNICSGNSESTQKCGIDVTL.
<b>Storage</b>	Lyophilized Osteoprotegerin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution OCIF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	TNFRSF11B tumor necrosis factor receptor superfamily, member 11b [ Homo sapiens ]
<b>Official Symbol</b>	TNFRSF11B
<b>Synonyms</b>	TNFRSF11B, OPG, OCIF, Osteoclastogenesis inhibitory factor, TR1, MGC29565,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Osteoprotegerin, TR1, Osteoclastogenesis inhibitory factor, tumor necrosis factor receptor superfamily, member 11b

**Gene ID**

4982

**mRNA Refseq**

NM\_002546

**Protein Refseq**

NP\_002537

**MIM**

602643

**UniProt ID**



O00300

**Chromosome  
Location**

8q24

**Pathway**Cytokine-cytokine receptor interaction; Monoamine Transport; Osteoclast Signaling;  
Osteoclast differentiation**Function**

cytokine activity; protein binding; receptor activity

 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA