

## Recombinant Human TNFRSF13B protein

Cat. No. TNFRSF13B-970H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human TNFRSF13B protein was expressed in Escherichia coli.
<b>Species</b>	Human
<b>Source</b>	E.coli
<b>ProteinLength</b>	160
<b>Description</b>	The protein encoded by this gene is a lymphocyte-specific member of the tumor necrosis factor (TNF) receptor superfamily. It interacts with calcium-modulator and cyclophilin ligand (CAML). The protein induces activation of the transcription factors NFAT, AP1, and NF-kappa-B and plays a crucial role in humoral immunity by interacting with a TNF ligand. This gene is located within the Smith-Magenis syndrome region on chromosome 17.
<b>Form</b>	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
<b>Bio-activity</b>	Fully biologically active when compared to standard. The biological activity is determined by its ability to block human BAFF induced T2B cell survival using a concentration range of 1.0-3.0 µg/ml.
<b>Molecular Mass</b>	Approximately 18.0 kDa, a single non-glycosylated polypeptide chain containing 160 amino acids.
<b>AA Sequence</b>	MSGGLGRSRRGGRSRVDQEERFPQGLWTGVAMRSCPEEQYWDPLLGTCSCKTIC

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

NHQSQR TCAAFCRSLSCRKEQGKFDHLLRDCISCASICGQHPKQCAYFCENKLRS  
PVNLPPELRRQRSGEVENNSDNSGRYQGLEHRGSEASPALPGLKLSADQV

**Endotoxin** Less than 1 EU/μg of rHuTACI/TNFRSF13B as determined by LAL method.

**Purity** >95% by SDS-PAGE and HPLC analysis.

**Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

**Reconstitution** We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

## GENE INFORMATION

**Gene Name** TNFRSF13B

**Official Symbol** TNFRSF13B

**Synonyms** TNFRSF13B; tumor necrosis factor receptor superfamily, member 13B; tumor necrosis factor receptor superfamily member 13B; CD267; TACI; tumor necrosis factor receptor 13B; transmembrane activator and CAML interactor; CVID; CVID2; TNFRSF14B; FLJ39942; MGC39952; MGC133214;

**Gene ID** 23495

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM\\_012452](#)

Protein Refseq [NP\\_036584](#)

MIM [604907](#)

UniProt ID [O14836](#)

**SDS-PAGE of  
TNFRSF13B-970H**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA