

# Active Recombinant Human TNFRSF17 Protein, Fc-tagged, FITC-Labeled

**Cat. No.** TNFRSF17-239H    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	FITC-Labeled Recombinant Human TNFRSF17 Protein (Met 1 - Ala 54) (Q02223), fused with a human IgG1 Fc tag at the C-terminus, was expressed in HEK293.
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Met 1 - Ala 54
<b>Form</b>	Lyophilized from sterile PBS, pH7.4, 10% trehalose.
<b>Bio-activity</b>	Measured by its binding ability in a functional ELISA. Immobilized Human BAFF, Fc Tag at 10µg/mL (100 µL/well) can bind FITC-Labeled Human BCMA, Fc Tag with a linear range of 0.15-0.625 µg/mL.
<b>Molecular Mass</b>	Recombinant Human TNFRSF17 Protein has a calculated MW of 32 kDa. As a result of glycosylation, The protein migrates as 38-42 kDa under reducing (R) condition and 60-70 kDa under non-reducing (NR) condition on SDS-PAGE gel.
<b>Endotoxin</b>	Less than 1.0 EU per µg by the LAL method.
<b>Purity</b>	>92% as determined by SDS-PAGE.
<b>Stability</b>	Samples are stable for up to twelve months from date of receipt at -70°C.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Storage</b>	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
<b>Conjugation</b>	FITC

## GENE INFORMATION

<b>Gene Name</b>	TNFRSF17 tumor necrosis factor receptor superfamily, member 17 [ Homo sapiens ]
<b>Official Symbol</b>	TNFRSF17
<b>Synonyms</b>	TNFRSF17; tumor necrosis factor receptor superfamily, member 17; BCMA; tumor necrosis factor receptor superfamily member 17; BCM; CD269; B-cell maturation factor; B cell maturation antigen; B-cell maturation protein;
<b>Gene ID</b>	608
<b>mRNA Refseq</b>	NM_001192
<b>Protein Refseq</b>	NP_001183
<b>MIM</b>	109545
<b>UniProt ID</b>	Q02223
<b>Chromosome Location</b>	16p13.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Pathway**

Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Intestinal immune network for IgA production, organism-specific biosystem; Intestinal immune network for IgA production, conserved biosystem;

**Function**

receptor activity; tumor necrosis factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA