

Active Recombinant Human TNFRSF9, MlgG2a Fc-tagged

Cat. No. TNFRSF9-234H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CD137 (AAH06196.1) (Glu19-Ser184) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell.
Species	Human
Source	CHO
ProteinLength	19-184 a.a.
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB.
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its binding ability in a functional ELISA.
Molecular Mass	44KDa (monomer)
AA Sequence	Glu19-Ser184

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="" test.="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	TNFRSF9 tumor necrosis factor receptor superfamily, member 9 [Homo sapiens]
Official Symbol	TNFRSF9
Synonyms	TNFRSF9; tumor necrosis factor receptor superfamily, member 9; ILA; tumor necrosis factor receptor superfamily member 9; 4 1BB; CD137; CD137 antigen; T cell antigen ILA; T-cell antigen ILA; 4-1BB ligand receptor; homolog of mouse 4-1BB; receptor protein 4-1BB; T-cell antigen 4-1BB homolog; induced by lymphocyte activation (ILA); interleukin-activated receptor, homolog of mouse Ly63; 4-1BB; CDw137; MGC2172; FLJ43501;
Gene ID	3604
mRNA Refseq	NM_001561
Protein Refseq	NP_001552
MIM	602250

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



UniProt ID	Q07011
Chromosome Location	1p36
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem;
Function	binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA