

Active Recombinant Mouse Tnfsf11 Protein, Fc-tagged, Alexa Fluor 647 conjugated

Cat. No. TNFSF11-860MAF647 Lot. No. (See product label)

SPECIFICATION

Product Overview	Alexa Fluor 647 conjugated recombinant extracellular domain of mouse TNFSF11 (BAA97257.1) (Arg 72-Asp 316) was fused with the Fc region of human IgG1 at the N-terminus.
Species	Mouse
Source	HEK293
ProteinLength	505
Form	Lyophilized
Bio-activity	Immobilized mouse Fc-TNFSF11 at 10 µg/mL (100 µL/well) can bind biotinylated human TNFRSF11B-His, The EC50 of biotinylated human TNFRSF11B-His is 0.07-0.17 µg/mL.
Molecular Mass	The recombinant mouse TNFSF11/Fc is a disulfide-linked homodimer. The reduced monomer comprises 505 amino acids and has a calculated molecular mass of 56 kDa. The apparent molecular mass of the monomer is approximately 50 kDa in SDS-PAGE under reducing conditions.
N-terminal Sequence Analysis	Glu

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	< 1.0 EU/ µg of the protein as determined by the LAL method.
Purity	> 87 % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer Labeled with Alexa Fluor 647 via amines Excitation = 650 nm Emission = 668 nm
Stability	Samples are stable for up to 12 months from date of receipt at -70 centigrade.
Storage	Store it under sterile conditions at -20 to-70 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Lyophilized from sterile PBS, pH 7.4, 5%-8% trehalose and mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
Conjugation	Alexa Fluor 647

GENE INFORMATION

Gene Name	Tnfsf11 tumor necrosis factor (ligand) superfamily, member 11 [<i>Mus musculus</i>]
Official Symbol	TNFSF11
Synonyms	TNFSF11; tumor necrosis factor (ligand) superfamily, member 11; tumor necrosis factor ligand superfamily member 11; OPG ligand; osteoprotegerin ligand; osteoclast differentiation factor; receptor activator of NF-kappaB ligand; TNF-related activation-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



	induc
Gene ID	21943
mRNA Refseq	NM_011613
Protein Refseq	NP_035743

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA