

## Recombinant Human TNFSF13 protein, Fc-tagged

Cat. No. TNFSF13-386H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human TNFSF13 protein (Ala105-Leu250), fused to human IgG1 Fc tag at N-terminus, was expressed in human 293 cells (HEK293).
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	146
<b>Description</b>	<p>APRIL(a proliferation-inducing ligand) is also known as Tumor necrosis factor ligand superfamily member 13, TALL-2, TRDL-1, CD256, TNFSF 13, cytokine that binds to TNFRSF13B/TACI and to TNFRSF17/BCMA. APRIL is a cytokine of the tumor necrosis factor family associated mainly with hematologic malignancies. The closely related TNF family ligands B cell activation factor (BAFF) and a proliferation-inducing ligand (APRIL) serve in the generation and maintenance of mature B-lymphocytes. Both BAFF and APRIL assemble as homotrimers that bind and activate several receptors that they partially share. BAFF-APRIL heteromers of different stoichiometries have distinct receptor-binding properties and activities. In addition, expression of APRIL was regulated by miR-145 in GC cells.</p>
<b>Form</b>	Lyophilized from 0.22 um filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
<b>Molecular Mass</b>	The protein has a calculated MW of 42.8 kDa. The protein migrates as 48 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Endotoxin</b>	Less than 1.0 EU per ug by the LAL method.
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Storage</b>	<p>For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower.</p> <p>Please avoid repeated freeze-thaw cycles.</p> <p>This product is stable after storage at:</p> <p>-20 centigrade to -70 centigrade for 12 months in lyophilized state;</p> <p>-70 centigrade for 3 months under sterile conditions after reconstitution.</p>
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

## GENE INFORMATION

<b>Gene Name</b>	TNFSF13
<b>Official Symbol</b>	TNFSF13
<b>Synonyms</b>	TNFSF13; tumor necrosis factor (ligand) superfamily, member 13; tumor necrosis factor ligand superfamily member 13; APRIL; CD256; a proliferation-inducing ligand; tumor necrosis factor-like protein ZTNF2; tumor necrosis factor-related death ligand-1; TNF- and APOL-related leukocyte expressed ligand 2; TALL2; ZTNF2; TALL-2; TRDL-1; FLJ57090; UNQ383/PRO715
<b>Gene ID</b>	8741
<b>mRNA Refseq</b>	NM_001198622
<b>Protein Refseq</b>	NP_001185551

 Tel: 1-631-559-9269 1-516-512-3133


 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**MIM** 604472


**UniProt ID** O75888

**SDS-PAGE of  
TNFSF13-386H**



Human APRIL, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

**Bioactivity-ELISA of  
TNFSF13-386H**




 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Immobilized Human BCMA, Mouse IgG2a Fc Tag, low endotoxin at 5 µg/mL (100 µL/well) can bind Human APRIL, Fc Tag with a linear range of 0.6-10 ng/mL (QC tested).

**Bioactivity-ELISA of  
TNFSF13-386H**



Immobilized Human APRIL, Fc Tag at 5 µg/mL (100 µL/well) can bind Biotinylated Human TACI, Fc, Avitag with a linear range of 0.01-0.313 µg/mL (Routinely tested).

☎ Tel: 1-631-559-9269 1-516-512-3133

☎ Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) ☎ Fax: 1-631-938-8127

☎ 45-1 Ramsey Road, Shirley, NY 11967, USA