

Recombinant Human TNFSF13B cell lysate

Cat. No. TNFSF13B-2025HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human BAFF / TNFSF13B derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

TNFSF13B tumor necrosis factor (ligand) superfamily, member 13b [Homo sapiens]

Official Symbol

TNFSF13B

Synonyms

TNFSF13B; tumor necrosis factor (ligand) superfamily, member 13b; TNFSF20; tumor necrosis factor ligand superfamily member 13B; BAFF; BLYS; CD257; TALL 1; TALL1; THANK; delta BAFF; Delta4 BAFF; B-lymphocyte stimulator; B-cell-activating factor; ApoL related ligand TALL-1; TNF homolog that activates apoptosis; dendritic cell-derived TNF-like molecule; tumor necrosis factor-like protein ZTNF4; TNF and ApoL-related leukocyte expressed ligand 1; tumor necrosis factor (ligand) superfamily, member 20; DTL; ZTNF4; TALL-1;

Gene ID

10673

mRNA Refseq

NM_001145645

Protein Refseq

NP_001139117

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	603969
UniProt ID	Q9Y275
Chromosome Location	13q32-q34
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Intestinal immune network for IgA production, organism-specific biosystem; Intestinal immune network for IgA production, conserved biosystem; Rheumatoid arthritis, organism-specific biosystem; Rheumatoid arthritis, conserved biosystem;
Function	cytokine activity; protein binding; receptor binding; tumor necrosis factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA