

## Recombinant Human TNFSF18, FLAG-tagged

Cat. No. TNFSF18-277H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	The extracellular domain of human GITRL (aa 50-177) is fused at the N-terminus to a FLAG-tag.
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	50-177 a.a.
<b>Description</b>	GITRL (Glucocorticoid-induced TNF receptor ligand) is expressed on dendritic cells (DC), monocytes, macrophages, B cells, activated T cells, endothelial cells, osteoclasts and various healthy non-lymphoid tissues (e.g. testis). GITRL is constitutively expressed and released as soluble form by solid tumors and various hematopoietic malignancies. GITRL causes differentiation of osteoclasts, activation of macrophages, but also alteration of carcinoma and leukemia cells and influences apoptosis. Binding to GITR is important in regulating T cell proliferation and TCR-mediated apoptosis. GITRL is implicated in development of autoimmune diseases and in the immune response against infectious pathogens and tumors.
<b>Form</b>	Lyophilized from 0.2µm-filtered solution in PBS.
<b>Molecular Mass</b>	~22kDa (SDS-PAGE)
<b>Purity</b>	≥90% (SDS-PAGE)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Stability</b>	Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Storage</b>	Short Term Storage: +4°C; Long Term Storage: -20°C. After opening, prepare aliquots and store at -20°C. Avoid freeze/thaw cycles.
<b>GENE INFORMATION</b>	
<b>Gene Name</b>	TNFSF18 tumor necrosis factor (ligand) superfamily, member 18 [ Homo sapiens ]
<b>Official Symbol</b>	TNFSF18
<b>Synonyms</b>	TNFSF18; tumor necrosis factor (ligand) superfamily, member 18; tumor necrosis factor ligand superfamily member 18; AITRL; hGITRL; TL6; AITR ligand; GITR ligand; activation-inducible TNF-related ligand; glucocorticoid-induced TNF-related ligand; glucocorticoid-induced TNFR-related protein ligand; GITRL; MGC138237;
<b>Gene ID</b>	<a href="#">8995</a>
<b>mRNA Refseq</b>	<a href="#">NM_005092</a>
<b>Protein Refseq</b>	<a href="#">NP_005083</a>
<b>MIM</b>	<a href="#">603898</a>
<b>UniProt ID</b>	<a href="#">Q9UNG2</a>
<b>Chromosome Location</b>	1q23
<b>Pathway</b>	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Function**

cytokine activity; receptor binding; tumor necrosis factor receptor binding; tumor necrosis factor receptor superfamily binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA