

Active Recombinant Human TPBG protein, His/Avi-tagged, Biotinylated

Cat. No. TPBG-051H Lot. No. (See product label)

SPECIFICATION

Product Overview

Biotinylated Recombinant Human TPBG(Ser31-Ser355) protein, fused to His/Avi tag at the C-terminus, was expressed in CHO cells .

Species

Human

Source

CHO

ProteinLength

Ser31-Ser355

Description

TPBG, also known as Trophoblast glycoprotein and Wnt-activated inhibitory factor 1 (WAIF1), is a single-pass transmembrane protein which may function as an inhibitor of Wnt/beta-catenin signaling (1). Distinct regions on the 5T4 ectodomain which contains eight leucine-rich repeats are thought to antagonize Wnt signaling (2). 5T4 has limited expression in normal adult tissues and in some solid tumors, while expression is elevated in human carcinomas which have been correlated with poor survival and prognosis (3-5). Human 5T4 is a 420 amino acid (aa) protein that includes a 31 aa signal peptide, a 324 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 44 aa cytoplasmic tail. Within the ECD, human 5T4 shares 78% and 77% aa sequence identity with mouse and rat 5T4, respectively. It has been identified that TIP-2/GIPC, a common interacting protein involved in cancer, interacts with 5T4 through its PDZ domain (6). 5T4 inhibits Wnt/ β -catenin signaling by binding to LRP6 and inhibit Wnt-induced LRP6 internalization into endocytic vesicles

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(1). Conversely, 5T4 can activate noncanonical Wnt signaling through DKK1 (1). In addition to LRP6, in-house data indicates 5T4 can also interact with Frizzled-8.

Predicted N Terminal Ser31

Form Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

Bio-activity Measured by its binding ability in a functional ELISA.

Molecular Mass 40-80 kDa, under reducing conditions

Endotoxin <0.10 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Applications Bioactivity

Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Reconstitution Reconstitute at 400 µg/mL in PBS.

Conjugation Biotin

GENE INFORMATION


Gene Name TPBG trophoblast glycoprotein [Homo sapiens]

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	TPBG
Synonyms	TPBG; trophoblast glycoprotein; 5T4; 5T4 AG; 5T4 oncofetal antigen; 5T4 oncotrophoblast glycoprotein; 5T4 oncofetal trophoblast glycoprotein; M6P1; 5T4AG;
Gene ID	7162
mRNA Refseq	NM_001166392
Protein Refseq	NP_001159864
MIM	190920
UniProt ID	Q13641

SDS-PAGE


2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Binding Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA