

Recombinant Human TRIB3 cell lysate

Cat. No. TRIB3-657HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human TRIB3 / NIPK derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

TRIB3 tribbles homolog 3 (Drosophila) [Homo sapiens]

Official Symbol

TRIB3

Synonyms

TRIB3; tribbles homolog 3 (Drosophila); C20orf97, chromosome 20 open reading frame 97; tribbles homolog 3; dJ1103G7.3; TRB3; TRB-3; p65-interacting inhibitor of NF-kappaB; p65-interacting inhibitor of NF-kappa-B; neuronal cell death inducible putative kinase; neuronal cell death-inducible putative kinase; NIPK; SINK; SKIP3; C20orf97;

Gene ID

57761

mRNA Refseq

NM_021158

Protein Refseq

NP_066981

MIM

607898

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q96RU7
Chromosome Location	20p13-p12.2
Pathway	Activation of PKB, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Adipogenesis, organism-specific biosystem; CD28 co-stimulation, organism-specific biosystem; CD28 dependent PI3K/Akt signaling, organism-specific biosystem; Costimulation by the CD28 family, organism-specific biosystem; Disease, organism-specific biosystem;
Function	ATP binding; enzyme binding; NOT kinase activity; protein binding; protein kinase binding; protein kinase binding; protein kinase inhibitor activity; transcription corepressor activity; ubiquitin protein ligase binding; ubiquitin-protein ligase regulator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA